

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number  
**WO 01/79187 A2**

(51) International Patent Classification: C07D 281/06,  
281/10, 417/04, 417/14, 513/04, 513/14, 405/04, A61K  
31/55, A61P 35/00

[US/US]; 8 Baycrest Way, South San Francisco, CA 94080  
(US).

(21) International Application Number: PCT/US01/12581

(74) Agents: ESMOND, Robert, W. et al.; Sterne, Kessler,  
Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Ave-  
nue, N.W., Washington, DC 20005-3934 (US).

(22) International Filing Date: 18 April 2001 (18.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/197,599 18 April 2000 (18.04.2000) US

(71) Applicant (for all designated States except US): CY-  
TOVIA, INC. [US/US]; 6650 Nancy Ridge Drive, San  
Diego, CA 92121 (US).

(81) Designated States (national): AE, AG, AI, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.


(71) Applicants and

(72) Inventors: CAI, Sui, Xiong [US/US]; 3623 Berryfield  
Court, San Diego, CA 92130 (US). DREWE, John, A.  
[US/US]; 7069 Murilla Lane, Carlsbad, CA 92009 (US).  
SHELTON, Emma, Jane [US/US]; 680 Lemon Street,  
Menlo Park, CA 94025 (US). LITVAK, Joane [US/US];  
555 Jean Street, Apartment 626, Oakland, CA 94610  
(US). SPERANDIO, David [CH/US]; 150 Paseo Court,  
Mountain View, CA 94043 (US). SPENCER, Jeffrey, R.



WO 01/79187 A2

(54) Title: SUBSTITUTED 1,4-THIAZEPINE AND ANALOGS AS ACTIVATORS OF CASPASES AND INDUCERS OF APOPTOSIS AND THE USE THEREOF



(I)

(57) Abstract: The present invention is directed to substituted 1,4-thiazepine and analogs thereof, represented by the general Formula (I): wherein the dashed lines, A¹, A², A³, X¹ and R¹ are defined herein. The present invention also relates to the discovery that compounds having Formula I are activators of caspases and inducers of apoptosis. Therefore, the activators of caspases and inducers of caspases and inducers of apoptosis of this invention can be used to induce cell death in a variety of clinical

conditions in which uncontrolled growth and spread of abnormal cells occurs.

# Substituted 1,4-Thiazepine and Analogs as Activators of Caspases and Inducers of Apoptosis and the Use Thereof

## Background of the Invention

### 5 *Field of the Invention*

This invention is in the field of medicinal chemistry. In particular, the invention relates to substituted 1,4-thiazepine and analogs, and the discovery that these compounds are activators of caspases and inducers of apoptosis. The invention also relates to the use of these compounds as therapeutically effective  
10 anti-cancer agents.

### *Related Art*

Organisms eliminate unwanted cells by a process variously known as regulated cell death, programmed cell death or apoptosis. Such cell death occurs as a normal aspect of animal development as well as in tissue homeostasis and aging (Glucksmann, A., *Biol. Rev. Cambridge Philos. Soc.* 26:59-86 (1951);  
15 Glucksmann, A., *Archives de Biologie* 76:419-437 (1965); Ellis, *et al.*, *Dev.* 112:591-603 (1991); Vaux, *et al.* *Cell* 76:777-779 (1994)). Apoptosis regulates cell number, facilitates morphogenesis, removes harmful or otherwise abnormal cells and eliminates cells that have already performed their function.  
20 Additionally, apoptosis occurs in response to various physiological stresses, such as hypoxia or ischemia (PCT published application WO96/20721).

There are a number of morphological changes shared by cells experiencing regulated cell death, including plasma and nuclear membrane

blebbing, cell shrinkage (condensation of nucleoplasm and cytoplasm), organelle relocalization and compaction, chromatin condensation and production of apoptotic bodies (membrane enclosed particles containing intracellular material) (Orrenius, S., *J. Internal Medicine* 237:529-536 (1995)).

5           Apoptosis is achieved through an endogenous mechanism of cellular suicide (Wyllie, A.H., in *Cell Death in Biology and Pathology*, Bowen and Lockshin, eds., Chapman and Hall (1991), pp. 9-34). A cell activates its internally encoded suicide program as a result of either internal or external signals. The suicide program is executed through the activation of a carefully  
10           regulated genetic program (Wyllie, *et al.*, *Int Rev. Cyt.* 68:251 (1980); Ellis, *et al.*, *Ann Rev. Cell Bio.* 7:663 (1991). Apoptotic cells and bodies are usually recognized and cleared by neighboring cells or macrophages before lysis. Because of this clearance mechanism, inflammation is not induced despite the clearance of great numbers of cells (Orrenius, S., *J Internal Medicine* 237:529-  
15           536 (1995)).

          A group of proteases are a key element in apoptosis (see, e.g., Thorneberry, *Chemistry and Biology* 5:R97-R103 (1998); Thornberry, *British Med. Bull.* 53:478-490 (1996)). Genetic studies in the nematode *Caenorhabditis elegans* revealed that apoptotic cell death involves at least 14 genes, two of  
20           which are the pro-apoptotic (death-promoting) *ced* (for *cell death abnormal*) genes, *ced-3* and *ced-4*. CED-3 is homologous to interleukin 1 beta-converting enzyme, a cysteine protease, which is now called caspase-1. Further extensive research revealed that the mammalian apoptosis system appears to involve a cascade of caspases, or a system that behaves like a cascade of caspases. At  
25           present, the caspase family of cysteine proteases comprises 14 different members, and more may be discovered in the future. All known caspases are synthesized as zymogens that require cleavage at an aspartyl residue prior to forming the active enzyme. Thus, caspases are capable of activating other caspases in the manner of an amplifying cascade.

Apoptosis and caspases are thought to be crucial in the development of cancer (*Apoptosis and Cancer Chemotherapy*, Hickman and Dive, eds., Humana Press (1999)). There is mounting evidence that cancer cells, while containing caspases, lack parts of the molecular machinery that activate the caspase cascade.

5 This makes the cancer cells lose their capacity to undergo cellular suicide and the cells become immortal, i.e., they become cancerous. Control points are known to exist in the apoptosis process that represent points for intervention leading to activation. These control points include the CED-9-BCL-like and CED-3-ICE-like gene family products, which are intrinsic proteins regulating the fate of a cell

10 to survive or die, respectively, and executing part of the cell death process itself (see, Schmitt, *et al.*, *Biochem. Cell. Biol.* 75:301-314 (1997)). BCL-like proteins include BCL-XL and BAX-alpha, which appear to function upstream of caspase activation. BCL-xL appears to prevent activation of the apoptotic protease cascade, whereas BAX-alpha accelerates activation of the apoptotic protease

15 cascade.

Chemotherapeutic (anti-cancer) drugs can trigger cancer cells to undergo suicide by activation of the dormant caspase cascade. This may be a crucial aspect of the mode of action of most, if not all, known anticancer drugs (Los, *et al.*, *Blood* 90:3118-3129 (1997); Friesen, *et al.*, *Nat. Med.* 2:574 (1996)). The

20 mechanism of action of current antineoplastic drugs frequently involves an attack at specific phases of the cell cycle. The cell cycle refers to the stages through which cells normally progress during their lifetimes. Normally, cells exist in a resting phase termed G<sub>0</sub>. During multiplication, cells progress to a stage in which DNA synthesis occurs, termed S. Later, cell division, or mitosis, occurs in a


25 phase called M. Antineoplastic drugs such as cytosine arabinoside, hydroxyurea, 6-mercaptopurine, and methotrexate are S phase specific, whereas antineoplastic drugs such as vincristine, vinblastine, and paclitaxel are M phase specific. Many slow growing tumors, for example colon cancers, exist primarily in the G<sub>0</sub> phase, whereas rapidly proliferating normal tissues, for example bone marrow, exist

30 primarily in the S or M phase. Thus, the possibility exists for the activation of the



caspase cascade, although the exact mechanisms for doing so presently are not clear. Furthermore, insufficient activity of the caspase cascade and consequent apoptotic events are implicated in various types of cancer. The development of caspase cascade activators and inducers of apoptosis is a highly desirable goal in the development of therapeutically effective antineoplastic agents. Moreover, since autoimmune disease and certain degenerative diseases also involve the proliferation of abnormal cells, therapeutic treatment for these diseases could be effected by enhancement of the apoptotic process through the administration of appropriate caspase cascade activators and inducers of apoptosis.

Sucheta *et al.*, (*Indian J. Chem. Sect. B: 34B:893-4 (1995)*) reported the synthesis of 1,4-benzothiazepine derivatives by reaction of 2-aminothiophenol with 3-cinnamoyl-2-pyrones ( $R = 2\text{-OMe}, 2\text{-}, 4\text{-OH}, 2\text{-}, 4\text{-NO}_2, 2,6\text{-Cl}_2, 2\text{-Cl}$ ):



### Summary of the Invention

The present invention is related to novel compounds of Formula I and the use of such compounds for treating, preventing or ameliorating neoplasia and cancer.

A second aspect of the present invention is related to the discovery that substituted 1,4-thiazepine and analogs are activators of the caspase cascade and inducers of apoptosis. Thus, an aspect of the present invention is directed to substituted 1,4-thiazepine and analogs as inducers of apoptosis.

A third aspect of the present invention is to provide a method for treating, preventing or ameliorating neoplasia and cancer by administering a compound of Formula I to a mammal in need of such treatment.


A fourth aspect of the present invention is to provide a pharmaceutical composition useful for treating disorders responsive to the induction of apoptosis, containing an effective amount of a compound of Formula I in admixture with one or more pharmaceutically acceptable carriers or diluents.

5 A fifth aspect of the present invention is directed to methods for the preparation of compounds of Formula I.

### *Detailed Description of the Preferred Embodiments*

The present invention arises out of the discovery that substituted 1,4-thiazepine and analogs, as represented in Formula I, are potent and highly efficacious activators of the caspase cascade and inducers of apoptosis. Therefore compounds of Formula I are useful for treating disorders responsive to induction of apoptosis.

Specifically, compounds useful in this aspect of the present invention are represented by Formula I:



in which:

the dashed lines indicate optional unsaturation without violating valency rules;

20  $R^1$  is hydrogen,  $(C_{1-6})$ alkyl or  $-C(O)R^6$ , wherein  $R^6$  is as defined below, or  $R^1$  is absent when a double bond exists between the nitrogen atom to which  $R^1$  is attached and an adjacent ring atom or  $R^1$  is as defined below;

$X^1$  is  $-NR^2$ ,  $-S$ ,  $-S(O)$ ,  $-S(O)_2$  or  $-O$ , wherein  $R^2$  is hydrogen or  $(C_{1-6})$ alkyl or is absent when a double bond exists between the nitrogen atom to which  $R^2$  is attached and an adjacent ring atom;

A<sup>1</sup> is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, or A<sup>1</sup> together with R<sup>1</sup> and the atoms to which A<sup>1</sup> and R<sup>1</sup> are attached forms a fused polycyclic ring system selected from heteroaryl and unsaturated, partially unsaturated or saturated heterocycloalkyl in any case containing a total of 10 to 15 ring atoms, wherein A<sup>1</sup> may be substituted with a group selected from -X<sup>2</sup>R<sup>3</sup>, -X<sup>2</sup>OR<sup>3</sup>, -X<sup>2</sup>C(O)R<sup>3</sup>, -X<sup>2</sup>OC(O)R<sup>3</sup>, -X<sup>2</sup>C(O)OR<sup>3</sup>, -X<sup>2</sup>SR<sup>3</sup>, -X<sup>2</sup>S(O)R<sup>3</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>3</sup>, -X<sup>2</sup>NR<sup>3</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>3</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>3</sup>, -X<sup>2</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>3</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>3</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene; R<sup>3</sup> is -X<sup>2</sup>R<sup>5</sup> wherein X<sup>2</sup> is as defined above and R<sup>5</sup> is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, wherein each ring within A<sup>1</sup> and R<sup>5</sup> contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within A<sup>1</sup> and R<sup>5</sup> may be substituted further with 1 to 2 groups independently selected from (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo, with the proviso that only one of A<sup>1</sup> and R<sup>5</sup> is a fused polycyclic ring system;


A<sup>2</sup> is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein A<sup>2</sup> may be substituted with a group selected

-7-

from  $-X^2R^8$ ,  $-X^2OR^8$ ,  $-X^2C(O)R^8$ ,  $-X^2OC(O)R^8$ ,  $-X^2C(O)OR^8$ ,  $-X^2SR^8$ ,  $-X^2S(O)R^8$ ,  $-X^2S(O)_2R^8$ ,  $-X^2NR^4R^8$ ,  $-X^2NR^4C(O)R^8$ ,  $-X^2NR^4C(O)OR^8$ ,  $-X^2C(O)NR^4R^8$ ,  $-X^2NR^4C(O)NR^4R^8$ ,  $-X^2NR^4C(NR^4)NR^4R^8$ ,  $-X^2NR^4S(O)_2R^8$  and  $-X^2S(O)_2NR^4R^8$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$  is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^2$  and  $R^8$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said heterocycloalkylene, carbocycloalkyl and heterocycloalkyl rings within  $A^2$  and  $R^8$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^2$  and  $R^8$  is a fused polycyclic ring system; and

$A^3$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-X^2R^9$ ,  $-X^2OR^9$ ,  $-X^2C(O)R^9$ ,  $-X^2OC(O)R^9$ ,  $-X^2C(O)OR^9$ ,  $-X^2SR^9$ ,  $-X^2S(O)R^9$ ,  $-X^2S(O)_2R^9$ ,  $-X^2NR^4R^9$ ,  $-X^2NR^4C(O)R^9$ ,  $-X^2NR^4C(O)OR^9$ ,  $-X^2C(O)NR^4R^9$ ,  $-X^2NR^4C(O)NR^4R^9$ ,  $-X^2NR^4C(NR^4)NR^4R^9$ ,  $-X^2NR^4S(O)_2R^9$  and  $-X^2S(O)_2NR^4R^9$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^9$  is  $-X^2R^{10}$  wherein  $X^2$  is as defined above and  $R^{10}$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10

ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^3$  and  $R^{10}$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within  $A^3$  and  $R^{10}$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^3$  and  $R^{10}$  is a fused polycyclic ring system; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof; with the proviso that when said compound is selected from the group consisting of Formulae II(a), II(b) and II(c):



then A<sup>3</sup> is other than:

unsubstituted pyridyl;

unsubstituted thienyl;

unsubstituted indolyl;

5 unsubstituted phenyl;


benzo[1,3]dioxolyl;

2,3-dihydro-benzo[1,4]dioxinyl;

phenyl which is mono-substituted by fluoro, bromo, iodo, methyl, isopropyl, ethoxy or methylsulfanyl; and

10 phenyl which is substituted by at least one of chloro, hydroxy or methoxy.

Alternatively, compounds useful for treating disorders responsive to induction of apoptosis are represented by Formula II:



in which:

15 the dashed lines indicate optional unsaturation without violating valency rules;

R<sup>1</sup> is hydrogen, (C<sub>1-6</sub>)alkyl or -C(O)R<sup>6</sup>, wherein R<sup>6</sup> is as defined below, or is absent when a double bond exists between the nitrogen atom to which R<sup>1</sup> is attached and an adjacent ring atom or R<sup>1</sup> is as defined below;

20 R<sup>7</sup> is hydrogen;

X<sup>1</sup> is -NR<sup>2</sup>-, -S-, -S(O)-, -S(O)<sub>2</sub>- or -O-, wherein R<sup>2</sup> is hydrogen or (C<sub>1-6</sub>)alkyl or is absent when a double bond exists between the nitrogen atom to which R<sup>2</sup> is attached and an adjacent ring atom;

25 A<sup>2</sup> is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a

total of 5 to 11 ring atoms, wherein  $A^2$  may be substituted with a group selected from  $-R^8$ ,  $-X^2OR^8$ ,  $-X^2C(O)R^8$ ,  $-X^2OC(O)R^8$ ,  $-X^2C(O)OR^8$ ,  $-X^2SR^8$ ,  $-X^2S(O)R^8$ ,  $-X^2S(O)_2R^8$ ,  $-X^2NR^4R^8$ ,  $-X^2NR^4C(O)R^8$ ,  $-X^2NR^4C(O)OR^8$ ,  $-X^2C(O)NR^4R^8$ ,  $-X^2NR^4C(O)NR^4R^8$ ,  $-X^2NR^4C(NR^4)NR^4R^8$ ,  $-X^2NR^4S(O)_2R^8$  and  $-X^2S(O)_2NR^4R^8$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$  is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^2$  and  $R^8$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said heteroalkylene, carbocycloalkyl and heterocycloalkyl rings within  $A^2$  and  $R^8$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo with the proviso that only one of  $A^2$  and  $R^8$  is a fused polycyclic ring system; and

$A^3$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-R^9$ ,  $-X^2OR^9$ ,  $-X^2C(O)R^9$ ,  $-X^2OC(O)R^9$ ,  $-X^2C(O)OR^9$ ,  $-X^2SR^9$ ,  $-X^2S(O)R^9$ ,  $-X^2S(O)_2R^9$ ,  $-X^2NR^4R^9$ ,  $-X^2NR^4C(O)R^9$ ,  $-X^2NR^4C(O)OR^9$ ,  $-X^2C(O)NR^4R^9$ ,  $-X^2NR^4C(O)NR^4R^9$ ,  $-X^2NR^4C(NR^4)NR^4R^9$ ,  $-X^2NR^4S(O)_2R^9$  and  $-X^2S(O)_2NR^4R^9$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^9$  is  $-X^2R^{10}$  wherein  $X^2$  is as defined above and  $R^{10}$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated,

partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^3$  and  $R^{10}$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within  $A^3$  and  $R^{10}$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo with the proviso that only one of  $A^3$  and  $R^{10}$  is a fused polycyclic ring system; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof;

provided, however, Formula II does not represent a compound wherein  $A^2$  is 2,3,6,7-tetrahydro-[1,4]thiazepinylene, 2,3-dihydrobenzo[*b*][1,4]thiazepinylene or 7-trifluoromethyl-2,3-dihydrobenzo[*b*][1,4]thiazepinylene when  $A^3$  is benzo[1,3]dioxolyl, indolyl, phenyl, pyridyl or thienyl, wherein said phenyl may be substituted with 1 to 3 groups independently selected from halo, nitro, hydroxy,  $(C_{1-4})$ alkyl,  $(C_{1-4})$ alkylsulfanyl and  $(C_{1-4})$ alkyloxy; or any *N*-oxide derivative, protected derivative, individual stereoisomer or mixture of stereoisomers, or pharmaceutically acceptable salt thereof.


Preferred are compounds of Formula I in which  $A^3$  is other than unsubstituted pyridyl; unsubstituted thienyl; unsubstituted indolyl; unsubstituted phenyl; benzo[1,3]dioxol-5-yl; 2,3-dihydro-benzo[1,4]dioxinyl; or phenyl which is substituted by at least one of halogen, nitro, hydroxy,  $(C_{1-3})$ alkyl, methoxy,



-12-


ethoxy and methylsulfanyl; and more preferred wherein A<sup>1</sup> is other than 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl.

Preferred are compounds of Formula I(A):




I(A)

wherein R<sup>1</sup>, A<sup>1</sup>, A<sup>2</sup> and A<sup>3</sup> are as in the Detailed Description of the Invention for Formula I; and more preferred are compounds of Formula I(B):



I(B)

wherein A<sup>1</sup>, A<sup>2</sup> and A<sup>3</sup> are as in the Detailed Description of the Invention for Formula I; and most preferred are compounds of Formula I in which A<sup>2</sup> is that is 2,3,6,7-tetrahydro-[1,4]thiazepin-5,7-ylene, that is a compound of Formula I(C):



I(C)

wherein A<sup>1</sup> and A<sup>3</sup> are as in the Detailed Description of the Invention for Formula I and said 2,3,6,7-tetrahydro-[1,4]thiazepin-5,7-ylene may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a

bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl.

Preferred are compounds of Formula I(C) in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and more preferred wherein said compound is selected from the group consisting of:

4-hydroxy-6-methyl-3-[7-(3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(5-ethyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(1-benzyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3[7-(2-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3[7-(3-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3[7-(4-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-[3-(3-trifluoromethyl-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-[3-(3,4-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-{7-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

3-{7-[5-(2-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[5-(3-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-[7-[5-(4-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-{7-[5-(chloro-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

5 3-[7-(4-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 3-[7-(1-benzenesulfonyl-1*H*-pyrrol-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(3-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

15 4-hydroxy-6-methyl-3-[7-(1-methyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(chloro-methyl-trifluoromethyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

20 3-{7-[1-(2,4-difluoro-benzenesulfonyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

25 3-{7-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-chloro-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

30 4-hydroxy-6-methyl-3-[7-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(4,5-dibromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methylsulfanyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

5 3-[7-(5-chloro-1-methyl-3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 4-hydroxy-6-methyl-3-[7-(4-trifluoromethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(bis-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-3-[7-(4-methanesulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; and

15 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-methoxy-6-methyl-pyran-2-one.

Preferred are compounds of Formula I(C) in which A<sup>1</sup> is A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl; and more preferred wherein said compound is selected from the group consisting of:

20 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

25 3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one; and

3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one.

-16-


Preferred are compounds of Formula I(C) in which A<sup>1</sup> is 2-hydroxy-6-oxo-cyclohex-1-enyl or 2-methoxy-6-oxo-cyclohex-1-enyl; and more preferred wherein said compound is selected from the group consisting of:

2-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

2-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone; and

3-hydroxy-2-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-cyclohex-2-enone.


Preferred are compounds of Formula I(C) in which A<sup>1</sup> is a group of Formula (c):



(c)

in which R<sup>7</sup> is hydrogen or methyl, R<sup>11</sup> is hydrogen or (C<sub>1-6</sub>)alkyl and the free valence is attached to A<sup>2</sup>; and in particular the compound of Formula I(C) which is 3-[7-2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-1H-quinolin-2-one.

Preferred are compounds of Formula I in which A<sup>2</sup> is 2,3-dihydro-benzo[b][1,4]thiazepin-5,7-ylene, that is a compound of Formula I(D):



I(D)

in which A<sup>1</sup> and A<sup>3</sup> are defined as in the Detailed Description of the Invention for Formula I, and said 2,3-dihydro-benzo[*b*][1,4]thiazepin-5,7-ylene may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>,  
 5 -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>,  
 -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>,  
 -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl.

10 Preferred are compounds of Formula I(D) in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and more preferred wherein said compound is 3-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-4-hydroxy-6-methyl-pyran-2-one.

15 Preferred are compounds of Formula I(D) in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl; and more preferred wherein said compound is 4-hydroxy-6-methyl-3-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-5,6-dihydro-pyran-2-one.


20 Preferred are compounds of Formula I(D) in which A<sup>1</sup> is 2-hydroxy-6-oxo-cyclohex-1-enyl or 2-methoxy-6-oxo-cyclohex-1-enyl; and more preferred wherein said compound is selected from the group consisting of:

3-hydroxy-2-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone; and

25 3-hydroxy-2-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone.

Preferred are compounds of Formula I in which A<sup>2</sup> is a group of Formula (k):

-18-



(k)

in which R<sup>1</sup> is defined as in the Detailed Description of the Invention for Formula I and said group of Formula (k) may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl; preferably wherein R<sup>1</sup> is hydrogen.

Preferred are compounds of Formula I in which in which A<sup>2</sup> is a group of Formula (k) and A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-5,6-dihydro-2H-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-5,6-dihydro-2H-pyran-3-yl; and more preferred wherein said compound is selected from the group consisting of:


3-[4-acetyl-7-(2,4-dimethoxy-phenyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one; and

3-[7-(2,4-dimethoxy-phenyl)-4-(2,2,2-trifluoro-ethanoyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one.

Preferred are compounds of Formula I in which in which A<sup>2</sup> is a group of Formula (k) and A<sup>1</sup> is optionally substituted phenyl; and more preferred wherein said compound is 1-[7-(2,4-dimethoxy-phenyl)-5-(3-fluoro-4-methoxyphenyl)-[1,4]thiazepan-4-yl]-ethanone.

Preferred are compounds of Formula I in which A<sup>2</sup> is 2,3-dihydro-[1,4]thiazepin-5,7-ylene, that is the compound of Formula I(F):

-19-



I(F)

in which A<sup>1</sup> and A<sup>3</sup> are defined as in Claim 1, and said 2,3-dihydro-[1,4]thiazepin-5,7-ylene may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl.

Preferred are compounds of Formula I(F) in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2H-pyran-3-yl; and more preferred wherein said compound is selected from the group consisting of:

3-[7-(2,4-dimethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one; and

3-(7-[2,2']bithienyl-5-yl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one.

Preferred are compounds of Formula I(F) in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-5,6-dihydro-2H-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-5,6-dihydro-2H-pyran-3-yl; and more preferred wherein said compound is 3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one.


Preferred are compounds of Formula I(F) in which A<sup>1</sup> is 2-hydroxy-6-oxo-cyclohex-1-enyl or 2-methoxy-6-oxo-cyclohex-1-enyl; and more preferred



-20-


wherein said compound is 2-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone.

Preferred are compounds of Formula I(G):



I(G)

5. wherein  $n$ ,  $R^1$ ,  $A^1$ ,  $A^2$  and  $A^3$  are as in the Detailed Description of the Invention for Formula I; and more preferred are compounds of Formula I(G) in which  $A^2$  is a group of Formula (I):



(I)


- 10 wherein  $n$  is 1 or 2; and said group of Formula (I) may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl.

- 15 Preferred are compounds of Formula I(G) in which  $A^2$  is a group of Formula (I) wherein  $n$  is 1 and  $A^1$  is 4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2H-pyran-3-yl; in particular the compound which is 20 3-[7-(2,4-dimethoxy-phenyl)-1-oxo-2,3,6,7-tetrahydro-1H-1λ⁴-[1,4]thiazepin-5-yl]-4-hydroxy-6-methoxy-pyran-2-one.

-21-


Preferred are compounds of Formula I(G) in which A<sup>2</sup> is a group of Formula (I) wherein n is 2 and A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2H-pyran-3-yl; in particular the compound which is 3-[7-(2,4-dimethoxy-phenyl)-1,1-dioxo-2,3,6,7-tetrahydro-1H-1λ<sup>6</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one.

Preferred are compounds of Formula I(I):



I(I)

wherein R<sup>1</sup>, R<sup>2</sup>, A<sup>1</sup>, A<sup>2</sup> and A<sup>3</sup> are as in the Detailed Description of the Invention for Formula I; and more preferred are compounds of Formula I(I) in which A<sup>2</sup> is a group of Formula (j):



(j)


wherein said group of Formula (I) may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl.

Preferred are compounds of Formula I(I) in which A<sup>2</sup> is a group of Formula (j) wherein A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl or 4-

-22-

methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; in particular the compound which is 3-[4-(2,4-dimethoxy-phenyl)-4,5-dihydro-3*H*-benzo[*b*][1,4]diazepin-2-yl]-4-hydroxy-6-methylpyran-2-one.

Preferred are compounds of Formula I(K):



I(K)

in which A<sup>3</sup> is defined as in the Detailed Description of the Invention for Formula I; and preferably the compound of Formula I(K) which is 10-(2,4-dimethoxy-phenyl)-3-methyl-7,8-dihydro-10*H*-2,5-dioxo-9-thia-6a-azacyclohepta[*a*]naphthalene-1,6-dione.

Preferred are compounds selected from the group consisting of:

4-hydroxy-3-[7-(2-methoxy-4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[7-(2-chloro-5-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;


3-[7-(4-dimethylamino-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

7-(2,4-dimethoxy-phenyl)-5-(4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl)-2,2-dimethyl-2,3,6,7-tetrahydro-[1,4]thiazepine-3-carboxylic acid; and


2-({1-[7-(2,4-dimethoxy-phenyl)-5-(4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl)-2,2-dimethyl-2,3,6,7-tetrahydro-[1,4]thiazepin-3-yl]-methanoyl}-amino)-propionic acid *tert*-butyl ester.

Preferred are compounds of Formula I in which A<sup>1</sup> is a group selected from Formulae (b), (c), (d), (e) and (f):


-23-




(b)




(c)



(d)




(e)



(f)

in which  $R^7$  is hydrogen or methyl,  $R^{11}$  is hydrogen or  $(C_{1-6})$ alkyl and the free valance is attached to  $A^2$ , or

$A^2$  and  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached form a group of Formula (g):



(g)


wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ ; and

$A^2$  is as defined above or is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein  $A^2$  may be


substituted with a group selected from  $-R^8$ ,  $-X^2OR^8$ ,  $-X^2C(O)R^8$ ,  $-X^2OC(O)R^8$ ,  $-X^2C(O)OR^8$ ,  $-X^2SR^8$ ,  $-X^2S(O)R^8$ ,  $-X^2S(O)_2R^8$ ,  $-X^2NR^4R^8$ ,  $-X^2NR^4C(O)R^8$ ,  $-X^2NR^4C(O)OR^8$ ,  $-X^2C(O)NR^4R^8$ ,  $-X^2NR^4C(O)NR^4R^8$ ,  $-X^2NR^4C(NR^4)NR^4R^8$ ,  $-X^2NR^4S(O)_2R^8$  and  $-X^2S(O)_2NR^4R^8$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$  is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^2$  and  $R^8$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said heteroalkylene, carbocycloalkyl and heterocycloalkyl rings within  $A^2$  and  $R^8$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^2$  and  $R^8$  is a fused polycyclic ring system.

Preferred are compounds of Formula I or II in which  $A^2$  is a group selected from Formulae (h), (i), (j), (k), (l) and (m):


-25-




(h)




(i)




(j)



(k)




(l)



(m)

in which  $n$  is 1 or 2 and  $R^1$  is acetyl or trifluoroacetyl or  $A^2$  and  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached form a group of Formula (g):



(g)

5 wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ .

Preferred are compounds of Formula I in which  $A^1$  is selected from Formulae (b) and (d), wherein  $R^7$  is hydrogen, and  $A^2$  is selected from Formulae (h) and (i).

Particularly preferred compounds of the invention include:

10 4-hydroxy-3-[7-(2-methoxy-4-methylsulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

4-hydroxy-3-[7-(4-methanesulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

2-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

10 3-hydroxy-2-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-cyclohex-2-enone; and

3-hydroxy-2-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone.

### ***Definitions:***

15 Unless otherwise stated, the following terms used in the specification and claims are defined for the purposes of this Application and have the following meanings.

"Alkyl" represented by itself means a straight or branched, saturated or unsaturated, aliphatic radical having the number of carbon atoms indicated (e.g., (C<sub>1-6</sub>)alkyl includes methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, isobutyl, 20 *tert*-butyl, vinyl, allyl, 1-propenyl, isopropenyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-methylallyl, ethynyl, 1-propynyl, 2-propynyl, and the like). Alkyl represented along with another radical (e.g., as in arylalkyl) means a straight or branched, saturated or unsaturated aliphatic divalent radical having the number of atoms indicated or when no atoms are indicated means a bond (e.g., (C<sub>6-10</sub>)aryl(C<sub>0-3</sub>)alkyl 25 includes phenyl, benzyl, phenethyl, 1-phenylethyl 3-phenylpropyl, and the like).

"Alkylene", unless indicated otherwise, means a straight or branched, saturated or unsaturated, aliphatic, divalent radical having the number of carbon atoms indicated (e.g., (C<sub>1-6</sub>)alkylene includes methylene (-CH<sub>2</sub>-), ethylene (-

$\text{CH}_2\text{CH}_2-$ ), trimethylene ( $-\text{CH}_2\text{CH}_2\text{CH}_2-$ ), tetramethylene ( $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ ), 2-butenylene ( $-\text{CH}_2\text{CH}=\text{CHCH}_2-$ ), 2-methyltetramethylene ( $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2-$ ), pentamethylene ( $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ ) and the like).

"Alkylidene" means a straight or branched saturated or unsaturated, aliphatic, divalent radical having the number of carbon atoms indicated (e.g. (C<sub>1-6</sub>)alkylidene includes methylene ( $=\text{CH}_2$ ), ethylidene ( $=\text{CHCH}_3$ ), isopropylidene ( $=\text{C}(\text{CH}_3)_2$ ), propylidene ( $=\text{CHCH}_2\text{CH}_3$ ), allylidene ( $=\text{CH}-\text{CH}=\text{CH}_2$ ), and the like).

"Amino" means the radical  $-\text{NH}_2$ . Unless indicated otherwise, the compounds of the invention containing amino moieties include protected derivatives thereof. Suitable protecting groups for amino moieties include acetyl, *tert*-butoxycarbonyl, benzyloxycarbonyl, and the like.

"Animal" includes humans, non-human mammals (e.g., dogs, cats, rabbits, cattle, horses, sheep, goats, swine, deer, and the like) and non-mammals (e.g., birds, and the like).

"Aromatic" means a moiety wherein the constituent atoms make up an unsaturated ring system, all atoms in the ring system are  $sp^2$  hybridized and the total number of pi electrons is equal to  $4n+2$ .

"Aryl" means an aromatic, monocyclic or fused bicyclic ring or ring assembly containing the total number of ring carbon atoms indicated, wherein each ring is comprised of 6 ring carbon atoms. Typical aryl groups containing a total of 6 to 14 ring atoms include phenyl, naphthyl, phenanthrenyl, anthracenyl, and the like. By further example, optionally substituted heteroaryl in defining A<sup>3</sup> includes 2,4-dimethoxyphenyl, 2-trifluoromethylsulfanylphenyl, 3-trifluoromethylsulfanylphenyl, 4-trifluoromethylsulfanylphenyl, 3-(3-trifluoromethylphenoxy)phenyl, 3-(3,4-dichlorophenoxy)phenyl, 3-(3,5-dichlorophenoxy)phenyl, 2-chloro-5-trifluoromethylphenyl, 2,4-difluorophenyl, 4-dimethylaminophenyl, 4-trifluoromethoxyphenyl, 2,4-di(trifluoromethyl)phenyl, 4-dimethylamino-2-methoxyphenyl, 4-



methylsulfonylphenyl, 2,4-diethoxyphenyl, 2,3,4-trimethoxyphenyl, 2-methoxy-4-methylsulfanyphenyl, and the like.

"Carbocycloalkyl" means a monocyclic, fused bicyclic or bridged polycyclic ring or ring assembly containing the number of ring carbon atoms indicated. "Unsaturated, partially unsaturated or saturated" used in connection with the term carbocycloalkyl refers to instances where the ring or ring assembly is unsaturated and non-aromatic (e.g., cyclooctatetraenyl and the like), partially saturated (e.g., azulenyl, fluorenyl, indenyl, 1,2,3,4-tetrahydronaphthyl, and the like) or saturated (e.g., cyclohexyl and the like), respectively.

"Disease" specifically includes any unhealthy condition of an animal or part thereof and includes an unhealthy condition that may be caused by, or incident to, medical or veterinary therapy applied to that animal, i.e., the "side effects" of such therapy.

"Halo" means fluoro, chloro, bromo or iodo.

"Halo-substituted alkyl", as an isolated group or part of a larger group, means "alkyl" substituted by one or more "halo" atoms, as such terms are defined in this Application. Halo-substituted alkyl includes haloalkyl, dihaloalkyl, trihaloalkyl, perhaloalkyl and the like (e.g. halo-substituted (C<sub>1-3</sub>)alkyl includes chloromethyl, dichloromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, perfluoroethyl, 2,2,2-trifluoro-1,1-dichloroethyl, and the like).

"Heteroatom moiety" includes -N=, -NR-, -O-, -S- or -S(O)<sub>2</sub>-, wherein R is hydrogen, (C<sub>1-6</sub>)alkyl or a protecting group.

"Heteroaryl" means an aromatic, monocyclic or fused bicyclic ring or ring assembly containing the total number of ring atoms indicated, wherein each ring is comprised of 5 or 6 ring atoms and one or more of the ring atoms is a heteroatom moiety selected from -N=, -NR-, -O- or -S-, wherein R is hydrogen, (C<sub>1-6</sub>)alkyl, a protecting group or represents the free valence serving as the point of attachment. For example, aryl containing a total of 5 to 14 ring atoms includes, but is not limited to acridine, benzo[b]thienyl, benzimidazolyl, benzoisoxazolyl, carbazolyl,  $\beta$ -carbolinyl, cinnolinyl, furazanyl, furyl, imidazolyl,

indolizinyl, 3*H*-indolyl, isobenzofuranyl, isoindolyl, isothiazolyl, isoquinolyl, isoxazolyl, naphtho[2,3]thienyl, naphthyridinyl, 2-oxobenzimidazolyl, perimidinyl, phenanthridinyl, phenazinyl, pteridinyl, purinyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, 2*H*-pyrrolyl, pyrrolyl, quinolyl, quinoxalyl, thienyl, and the like. By further example, optionally substituted heteroaryl in defining A<sup>3</sup> includes 3-phenyl-1*H*-pyrazol-4-yl, 5-ethylthien-2-yl, 1-benzyl-1*H*-indol-3-yl, 5-(3-trifluoromethylphenyl)fur-2-yl, 5-(2-chlorophenyl)fur-2-yl, 5-(3-chlorophenyl)fur-2-yl, 5-(4-chlorophenyl)fur-2-yl, 5-(2-chloro-5-trifluoromethylphenyl)fur-2-yl, 4-bromothien-2-yl, 5-bromothien-2-yl, 1-phenylsulfonyl-1*H*-pyrrol-2-yl, 3-methylfur-2-yl, 5-methylfur-2-yl, 1-methyl-1*H*-indol-3-yl, 5-chloro-1-methyl-3-trifluoromethyl-1*H*-pyrrol-5-yl, 4[2,2']bithienyl-5-yl, 1-(3,5-dichloro)pyrrol-2-yl, 1-(4-chloro)pyrrol-2-yl, 5-chloro-1*H*-indol-3-yl, 6-(4-methylphenylsulfonyl)imidazo[2,1-*b*]thiazol-5-yl, 4,5-dibromothien-2-yl, 5-methylsulfonylthien-2-yl, 5-chloro-1-methyl-3-phenyl-1*H*-pyrrol-4-yl, and the like.

"Heteroarylene" means a divalent, aromatic, monocyclic or fused bicyclic ring or ring assembly containing the total number of ring atoms indicated, wherein each ring is comprised of 5 or 6 ring atoms and two or more of the ring atoms are a heteroatom moiety selected from -N=, -NR-, -O- or -S-, wherein R is hydrogen, (C<sub>1-6</sub>)alkyl or a protecting group. For example, aryl containing a total of 5 to 14 ring atoms includes, but is not limited to pyrimidin-2,4-ylene, pyrrolo[1,2-*a*]pyrimidin-2,4-ylene, and the like.

"Heterocycloalkylene" means a divalent, monocyclic, fused bicyclic or bridged polycyclic ring or ring assembly containing the number of ring atoms indicated, wherein two or more of the ring atoms are a heteroatom moiety selected from -N=, -NR-, -O- or -S-, wherein R is hydrogen, (C<sub>1-6</sub>)alkyl or a protecting group. "Unsaturated, partially unsaturated or saturated" used in connection with the term heterocycloalkylene refers to instances where the ring or ring assembly is unsaturated and non-aromatic (e.g., 1,4-thiazepin-5,7-ylene and the like), partially saturated (e.g., 2,3-dihydro-1,4-thiazepin-5,7-ylene, 2,3-

dihydrobenzo[*b*][1,4]thiazepin-5,7-ylene, 2,3-dihydrobenzo[*b*][1,4]diazepin-5,7-ylene, and the like) or saturated (e.g. 1,4-thiazepan-5,7-ylene and the like), respectively. For example, optionally substituted heterocycloalkylene used in defining A<sup>2</sup> includes 2,3,6,7-tetrahydro[1,4]thiazepin-5,7-ylene, 2,3-dihydrobenzo[*b*][1,4]thiazepin-5,7-ylene, 4-acetyl[1,4]thiazepan-5,7-ylene, 4-trifluoroacetyl[1,4]thiazepan-5,7-ylene, 2,3-dihydrobenzo[*b*][1,4]diazepin-5,7-ylene, 2,3-dihydro-1,4-thiazepin-5,7-ylene, 1-oxo-2,3,6,7-tetrahydro[1,4]thiazepin-5,7-ylene, 1,1-dioxo-2,3,6,7-tetrahydro[1,4]thiazepin-5,7-ylene, 3-(1-*tert*-butoxycarbonyl ethyl carbamoyl)-2,3,6,7-tetrahydro[1,4]thiazepin-5,7-ylene, and the like.

"Heterocycloalkyl" means a monocyclic, fused bicyclic or bridged polycyclic ring or ring assembly containing the number of ring atoms indicated, wherein one or more of the ring atoms is a heteroatom moiety selected from -N=, -NR-, -O- or -S-, wherein R is hydrogen, (C<sub>1-6</sub>)alkyl or a protecting group.

"Unsaturated, partially unsaturated or saturated" used in connection with the term heterocycloalkyl refers to instances where the ring or ring assembly is unsaturated and non-aromatic (e.g., oxepinyl, thiepinyl, and the like), partially saturated (e.g., 2*H*-pyranyl, 3,6-dihydro-2*H*-pyran, 1,2-dihydroquinolyl, and the like) or saturated (e.g. tetrahydropyranyl, and the like), respectively. By further example, optionally substituted heterocycloalkyl used in defining A<sup>1</sup> includes 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl, 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl, 4-hydroxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl, 2-hydroxy-6-oxo-cyclohex-1-enyl, 4-hydroxy-2-oxo-1,2-dihydro-quinolin-3-yl, 7-amino-1-oxo-1*H*-isochromen-8-yl, 2,3-dioxo-3,4-dihydro-2*H*-quinoxalin-1-yl, 2-oxo-2,3-dihydro-indol-1-yl, 4-oxo-4*H*-pyrido[1,2-*a*]pyrimidin-3-yl, and the like.

"Imino" means the moiety =NR, wherein R is hydrogen or (C<sub>1-6</sub>)alkyl.

"Isomers" mean compounds of Formula I having identical molecular formulae but differ in the nature or sequence of bonding of their atoms or in the arrangement of their atoms in space. Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not

mirror images of one another are termed "diastereomers" and stereoisomers that are nonsuperimposable mirror images are termed "enantiomers" or sometimes "optical isomers". A carbon atom bonded to four nonidentical substituents is termed a "chiral center". A compound with one chiral center has two enantiomeric forms of opposite chirality is termed a "racemic mixture". A compound that has more than one chiral center has  $2^{n-1}$  enantiomeric pairs, where  $n$  is the number of chiral centers. Compounds with more than one chiral center may exist as either an individual diastereomers or as a mixture of diastereomers, termed a "diastereomeric mixture". When one chiral center is present a stereoisomer may be characterized by the absolute configuration of that chiral center. Absolute configuration refers to the arrangement in space of the substituents attached to the chiral center. Enantiomers are characterized by the absolute configuration of their chiral centers and described by the *R*- and *S*-sequencing rules of Cahn, Ingold and Prelog. Conventions for stereochemical nomenclature, methods for the determination of stereochemistry and the separation of stereoisomers are well known in the art (e.g., see "Advanced Organic Chemistry", 4th edition, March, Jerry, John Wiley & Sons, New York, 1992). It is understood that the names and illustration used in this Application to describe compounds of Formula I are meant to be encompassed all possible stereoisomers.

"Nitro" means the radical  $-\text{NO}_2$ .

"Oxo" means the moiety  $=\text{O}$ .

"Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not.

"*N*-Oxide derivatives" means derivatives of compounds of Formula I in which nitrogens are in an oxidized state (i.e., O-N) and which possess the desired pharmacological activity.

"Pathology" of a disease means the essential nature, causes and development of the disease as well as the structural and functional changes that result from the disease processes.

"Pharmaceutically acceptable" means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary use as well as human pharmaceutical use.

"Pharmaceutically acceptable salts" means salts of compounds of Formula I which are pharmaceutically acceptable, as defined above, and which possess the desired pharmacological activity. Such salts include acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or with organic acids such as acetic acid, propionic acid, hexanoic acid, heptanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartatic acid, citric acid, benzoic acid, *o*-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, maleic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, *p*-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, *p*-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid and the like.

Pharmaceutically acceptable salts also include base addition salts which may be formed when acidic protons present are capable of reacting with inorganic or organic bases. Acceptable inorganic bases include sodium hydroxide, sodium carbonate, potassium hydroxide, aluminum hydroxide and calcium hydroxide. Acceptable organic bases include ethanolamine, diethanolamine, triethanolamine, tromethamine, *N*-methylglucamine and the like.

"Prodrug" means a compound which is convertible in vivo by metabolic means (e.g. by hydrolysis) to a compound of the invention. For example an ester of a compound of the invention containing a hydroxy group may be convertible by hydrolysis in vivo to the parent molecule. Alternatively an ester of a compound of the invention containing a carboxy group may be convertible by hydrolysis in vivo to the parent molecule. Suitable esters of compounds of the invention containing a hydroxy group, are for example acetates, citrates, lactates, tartrates, malonates, oxalates, salicylates, propionates, succinates, fumarates, maleates, methylene-bis-b-hydroxynaphthoates, gentisates, isethionates, di-*p*-toluoyltartrates, methanesulphonates, ethanesulphonates, benzenesulphonates, *p*-toluenesulphonates, cyclohexylsulphamates and quinate. Suitable esters of compounds of the invention containing a carboxy group, are for example those described by F.J. Leinweber, *Drug Metab. Res.*, 1987, 18, page 379. An especially useful class of esters of compounds of the invention containing a hydroxy group, may be formed from acid moieties selected from those described by Bundgaard et al., *J. Med. Chem.*, 1989, 32, page 2503-2507, and include substituted (aminomethyl)-benzoates, for example, dialkylaminomethylbenzoates in which the two alkyl groups may be joined together and/or interrupted by an oxygen atom or by an optionally substituted nitrogen atom, e.g. an alkylated nitrogen atom, more especially (morpholino-methyl)benzoates, e.g. 3- or 4-(morpholinomethyl)-benzoates, and (4-alkylpiperazin-1-yl)benzoates, e.g. 3- or 4-(4-alkylpiperazin-1-yl)benzoates.

"Protected derivatives" means derivatives of compounds of the invention in which a reactive site or sites are blocked with protecting groups. Protected derivatives of compounds of the invention are useful in the preparation of compounds of the invention or in themselves may be active as activators of the caspase cascade and inducers of apoptosis. A comprehensive list of suitable protecting groups can be found in T.W. Greene, *Protecting Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons, Inc. 1999.

"Therapeutically effective amount" means that amount which, when administered to an animal for treating a disease, is sufficient to effect such treatment for the disease.

"Thioxo" means the moiety =S.

5 "Treatment" or "treating" means any administration of a compound of the present invention and includes:

(1) preventing the disease from occurring in an animal which may be predisposed to the disease but does not yet experience or display the pathology or symptomatology of the disease,

10 (2) inhibiting the disease in an animal that is experiencing or displaying the pathology or symptomatology of the disease (i.e., arresting further development of the pathology and/or symptomatology), or

(3) ameliorating the disease in an animal that is experiencing or displaying the pathology or symptomatology of the disease (i.e., reversing the pathology and/or symptomatology).

#### *Nomenclature:*

The compounds of Formulae I and II and the intermediates and starting materials used in their preparation are named in accordance with IUPAC rules of nomenclature in which the characteristic groups have decreasing priority for citation as the principle group, e.g., acids, esters, amides, etc, as determined by  
20 AutoNom 4.0 (Beilstein Information Systems, Inc.). For example, a compound of Formula II wherein A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxopyran-3-yl, A<sup>2</sup> is 2,3,6,7-tetrahydro[1,4]thiazepin-5,7-ylene and A<sup>3</sup> is 3-(3,5-dichloro-phenoxy)-phenyl is named 3-{7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-  
25 5-yl}-4-hydroxy-6-methyl-pyran-2-one. Alternatively, the compound may be referred to as 5-(4-hydroxy-6-methyl-2H-pyran-2-one-3-yl)-7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepine.

**Utility and Pharmacology:**

Another important aspect of the present invention is the discovery that substituted 1,4-thiazepines and analogs thereof are potent and highly efficacious activators of caspases and inducers of apoptosis in drug resistant cancer cells, such as breast and prostate cancer cells, which enables these compounds to kill these drug resistant cancer cells. In comparison, most standard anti-cancer drugs are not effective in killing drug resistant cancer cells under the same conditions. Therefore, substituted 1,4-thiazepines and analogs thereof are useful for the treatment of drug resistant cancer in animals.

The present invention includes a therapeutic method useful to modulate *in vivo* apoptosis or *in vivo* neoplastic disease, comprising administering to a subject in need of such treatment an effective amount of a substituted 1,4-thiazepines or analog thereof, or a pharmaceutically acceptable salt or prodrug thereof, which functions as a caspase cascade activator and inducer of apoptosis.

The present invention also includes a therapeutic method comprising administering to an animal an effective amount of a substituted 1,4-thiazepines or analog thereof, or a pharmaceutically acceptable salt or prodrug thereof, wherein said therapeutic method is useful to treat cancer, which is a group of diseases characterized by the uncontrolled growth and spread of abnormal cells. Such diseases include, but are not limited to, Hodgkin's disease, non-Hodgkin's lymphomas, acute and chronic lymphocytic leukemias, multiple myeloma, neuroblastoma, breast carcinomas, ovarian carcinomas, lung carcinomas, Wilms' tumor, cervical carcinomas, testicular carcinomas, soft tissue sarcomas, chronic lymphocytic leukemia, primary macroglobulinemia, bladder carcinomas, chronic granulocytic leukemia, primary brain carcinomas, malignant melanoma, small-cell lung carcinomas, stomach carcinomas, colon carcinomas, malignant pancreatic insulinoma, malignant carcinoid carcinomas, malignant melanomas, choriocarcinomas, mycosis fungoides, head and neck carcinomas, osteogenic sarcoma, pancreatic carcinomas, acute granulocytic leukemia, hairy cell leukemia,



neuroblastoma, rhabdomyo sarcoma, Kaposi's sarcoma, genitourinary carcinomas, thyroid carcinomas, esophageal carcinomas, malignant hypercalcemia, cervical hyperplasia, renal cell carcinomas, endometrial carcinomas, polycythemia vera, essential thrombocytosis, adrenal cortex carcinomas, skin cancer, and prostatic carcinomas.

In practicing the therapeutic methods, effective amounts of compositions containing therapeutically effective concentrations of the compounds formulated for oral, intravenous, local and topical application, for the treatment of neoplastic diseases and other diseases in which caspase cascade mediated physiological responses are implicated, are administered to an individual exhibiting the symptoms of one or more of these disorders. The amounts are effective to ameliorate or eliminate one or more symptoms of the disorders. An effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce the symptoms associated with the disease. Such amount may be administered as a single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but, typically, is administered in order to ameliorate the disease. Typically, repeated administration is required to achieve the desired amelioration of symptoms.

In another embodiment, a pharmaceutical composition comprising a substituted 1,4-thiazepines or analog thereof, or a pharmaceutically acceptable salt thereof, which functions as a caspase cascade activator and inducer of apoptosis in combination with a pharmaceutically acceptable vehicle is provided.

Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a substituted 1,4-thiazepines or analog thereof, or a pharmaceutically acceptable salt or prodrug thereof, which functions as a caspase cascade activator and inducer of apoptosis, in combination with at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent. Examples of known anticancer agents which can be used for combination therapy include, but not are limit to alkylating agents such as

busulfan, cis-platin, mitomycin C, and carboplatin; antimitotic agents such as colchicine, vinblastine, paclitaxel, and docetaxel; topo I inhibitors such as camptothecin and topotecan; topo II inhibitors such as doxorubicin and etoposide; RNA/DNA antimetabolites such as 5-azacytidine, 5-fluorouracil and methotrexate; DNA antimetabolites such as 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea and thioguanine; and antibodies such as Herceptin® and Rituxan®. Other known anti-cancer agents which can be used for combination therapy include melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen and alanosine.

In practicing the methods of the present invention, the substituted 1,4-thiazepines or analog thereof may be administered together with at least one known chemotherapeutic agent as part of a unitary pharmaceutical composition. Alternatively, the compound of the invention may be administered apart from at least one known cancer chemotherapeutic agent. In one embodiment, the compound of the invention and at least one known cancer chemotherapeutic agent are administered substantially simultaneously, i.e. the compounds are administered at the same time or one after the other, so long as the compounds reach therapeutic levels in the blood at the same time. In another embodiment, the compound of the invention and at least one known cancer chemotherapeutic agent are administered according to their individual dose schedule, so long as the compounds reach therapeutic levels in the blood.

Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a bioconjugates of the substituted 1,4-thiazepines or analogs thereof, which functions as a caspase cascade activator and inducer of apoptosis, in bioconjugation with at least one known therapeutically useful antibody, such as Herceptin® or Rituxan®, growth factor such as DGF or NGF, cytokines such as IL-2 or IL-4, or any other molecule that binds to a cell surface receptor. These conjugates can be made using functional groups of the substituted 1,4-thiazepines or analogs thereof. For example, when one of the R

groups is a carboxylic acid, it can be used to modify the amino group on the proteins, to produce a drug-protein conjugate. For example, the *N*-hydroxysuccinamide ester of the carboxy-containing compound may be prepared which may be condensed with the protein. See *Anal. Biochem* 87:218 (1994) and  
5 *Polycyclic Aromat. Compd* 3:781 (1993). The antibodies and other molecules will deliver the substituted 1,4-thiazepine or analog thereof to its targets and make them more effective anticancer agents. The bioconjugates also may enhance the anticancer effect of therapeutically useful antibodies, such as Herceptin® or Rituxan®.

10 Similarly, another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a substituted 1,4-thiazepine or analog thereof, or a pharmaceutically acceptable salt or prodrug thereof, which functions as a caspase cascade activator and inducer of apoptosis, in combination with radiation therapy. In this embodiment, the substituted 1,4-thiazepine or  
15 analog thereof may be administered at the same time as the radiation therapy is administered or at a different time.

Yet another embodiment of the present invention is directed to a composition effective for post-surgical treatment of cancer, comprising a substituted 1,4-thiazepine or analog thereof, or a pharmaceutically acceptable salt  
20 or prodrug thereof, which functions as a caspase cascade activator and inducer of apoptosis. The invention also relates to a method of treating cancer by surgically removing the cancer and then treating the animal with one of the pharmaceutical compositions described herein.

A wide range of immune mechanisms operate rapidly following exposure  
25 to an infectious agent. Depending on the type of infection, rapid clonal expansion of the T and B lymphocytes occurs to combat the infection. The elimination of the effector cells following an infection is one of the major mechanisms maintaining immune homeostasis. This deletion of reactive cell has been shown to be regulated by a phenomenon known as apoptosis. Autoimmune diseases  
30 have been lately identified as a consequence of deregulated cell death. In certain

autoimmune diseases, the immune system directs its powerful cytotoxic effector mechanisms against specialized cells such as oligodendrocytes in multiple sclerosis, the beta cells of the pancreas in diabetes mellitus, and thyrocytes in Hashimoto's thyroiditis (Ohsako, S. & Elkon, K.B., *Cell Death Differ.* 6:13-21 (1999)). Mutations of the gene encoding the lymphocyte apoptosis receptor Fas/APO-1/CD95 are reported to be associated with defective lymphocyte apoptosis and autoimmune lymphoproliferative syndrome (ALPS), which is characterized by chronic, histologically benign splenomegaly and generalized lymphadenopathy, hypergammaglobulinemia, and autoantibody formation (Infante, A.J., et al., *J Pediatr.* 133:629-633 (1998) and Vaishnaw, A.K., et al., *J Clin. Invest.* 103:355-363 (1999)). Overexpression of Bcl-2, which is a member of the bcl-2 gene family of programmed cell death regulators with anti-apoptotic activity in developing B cells of transgenic mice, in the presence of T cell dependent co-stimulatory signals, results in the generation of a modified B cell repertoire and in the production of pathogenic autoantibodies (Lopez-Hoyos, M., et al., *Int. J. Mol. Med.* 1:475-483 (1998)). Accordingly, many types of autoimmune disease may be caused by defects of the apoptotic process, and one treatment strategy would be to turn on apoptosis in the lymphocytes that are causing autoimmune disease (O'Reilly, L.A. & Strasser, A., *Inflamm. Res.* 48:5-21 (1999)).

Fas-Fas ligand (FasL) interaction is known to be required for the maintenance of immune homeostasis. Experimental autoimmune thyroiditis (EAT), characterized by autoreactive T and B cell responses and a marked lymphocytic infiltration of the thyroid, is a good model to study the therapeutic effects of FasL. Batteux, F., et al., (*J Immunol.* 162:603-608 (1999)) reported that by direct injection of DNA expression vectors encoding FasL into the inflamed thyroid, the development of lymphocytic infiltration of the thyroid was inhibited and induction of infiltrating T cells death was observed. These results show that FasL expression on thyrocytes may have a curative effect on ongoing EAT by inducing death of pathogenic autoreactive infiltrating T lymphocytes.

Bisindolylmaleimide VIII is known to potentiate Fas-mediated apoptosis in human astrocytoma 1321NI cells and in Molt-4T cells, and both of which were resistant to apoptosis induced by anti-Fas antibody in the absence of bisindolylmaleimide VIII. Potentiation of Fas-mediated apoptosis by bisindolylmaleimide VIII was reported to be selective for activated, rather than non-activated, T cells, and was Fas-dependent. Zhou T., *et al.*, (*Nat. Med* 5:42-49 (1999)) reported that administration of bisindolylmaleimide VIII to rats during autoantigen stimulation prevented the development of symptoms of T cell-mediated autoimmune diseases in two models, the Lewis rat model of experimental allergic encephalitis and the Lewis adjuvant arthritis model. Therefore the application of a Fas-dependent apoptosis enhancer such as bisindolylmaleimide VIII may be therapeutically useful for the more effective elimination of detrimental cells and inhibition of T cell-mediated autoimmune diseases. Therefore an effective amount of a substituted 1,4-thiazepines or analog thereof, or a pharmaceutically acceptable salt or prodrug thereof, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for autoimmune disease.

Psoriasis is a chronic skin disease that is characterized by scaly red patches. Psoralen plus ultraviolet A (PUVA) is a widely used and effective treatment for psoriasis vulgaris and Coven, *et al.*, *Photodermatol. Photoimmunol. Photomed* 15:22-27 (1999), reported that lymphocytes treated with psoralen 8-MOP or TMP plus UVA displayed DNA degradation patterns typical of apoptotic cell death. Ozawa, *et al.*, *J Exp. Med* 189:711-718 (1999) reported that induction of T cell apoptosis could be the main mechanism by which 312-nm UVB resolves psoriasis skin lesions. Low doses of methotrexate may be used to treat psoriasis to restore a clinically normal skin. Heenen, *et al.*, *Arch. Dermatol. Res.* 290:240-245 (1998), reported that low doses of methotrexate may induce apoptosis and this mode of action could explain the reduction in epidermal hyperplasia during treatment of psoriasis with methotrexate. Therefore an effective amount of a substituted 1,4-thiazepine or analog thereof, or a pharmaceutically acceptable salt

or prodrug thereof, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for psoriasis.

Synovial cell hyperplasia is a characteristic of patients with rheumatoid arthritis (RA). Excessive proliferation of RA synovial cells as well as defects in synovial cell death might be responsible for the synovial cell hyperplasia. Wakisaka, *et al.*, *Clin. Exp. Immunol.* 114:119-128 (1998), found that although RA synovial cells could die via apoptosis through Fas/FasL pathway, apoptosis of synovial cells was inhibited by proinflammatory cytokines present within the synovium, and suggested that inhibition of apoptosis by the proinflammatory cytokines may contribute to the outgrowth of synovial cells, and lead to pannus formation and the destruction of joints in patients with RA. Therefore an effective amount of a substituted 1,4-thiazepine or analog thereof, or a pharmaceutically acceptable salt or prodrug thereof, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for rheumatoid arthritis.

An accumulation of convincing evidence suggests that apoptosis plays a major role in promoting resolution of the acute inflammatory response. Neutrophils are constitutively programmed to undergo apoptosis, thus limiting their pro-inflammatory potential and leading to rapid, specific, and non-phlogistic recognition by macrophages and semi-professional phagocytes (Savill, J., *J Leukoc. Biol.* 61:375-380 (1997)). Boirivant, *et al.*, *Gastroenterology* 116:557-565 (1999), reported that lamina propria T cells isolated from areas of inflammation in Crohn's disease, ulcerative colitis, and other inflammatory states manifest decreased CD2 pathway-induced apoptosis, and that studies of cells from inflamed Crohn's disease tissue indicate that this defect is accompanied by elevated Bel-2 levels. Therefore an effective amount of a substituted 1,4-thiazepines or analog thereof, or a pharmaceutically acceptable salt or prodrug thereof, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for inflammation and inflammatory bowel disease.

Compositions within the scope of this invention include all compositions wherein the caspase cascade activators are contained in an amount that is effective to achieve its intended purpose. While individual needs vary, determination of optimal ranges of effective amounts of each component is within the skill of the art. Typically, the active compounds may be administered to mammals, e.g., humans, orally at a dose of 0.0025 to 50 mg/kg, or an equivalent amount of the pharmaceutically acceptable salt thereof, per day of the body weight of the mammal being treated for apoptosis mediated disorders. Preferably, about 0.01 to about 10 mg/kg is orally administered to treat or prevent such disorders. For intramuscular injection, the dose is generally about one-half of the oral dose. For example, a suitable intramuscular dose would be about 0.0025 to about 25 mg/kg, and most preferably, from about 0.01 to about 5 mg/kg. If a known cancer chemotherapeutic agent is also administered, it is administered in an amount with is effective to achieve its intended purpose. The amounts of such known cancer chemotherapeutic agents effective for cancer are well known to those of skill in the art.

The unit oral dose may comprise from about 0.01 to about 50 mg, preferably about 0.1 to about 10 mg of the caspase cascade activator. The unit dose may be administered one or more times daily as one or more tablets each containing from about 0.1 to about 10, conveniently about 0.25 to 50 mg of the compound or its solvates. In a topical formulation, the caspase cascade activator may be present at a concentration of about 0.01 to 100 mg per gram of carrier.

In addition to administering the caspase cascade activator as a raw chemical, the caspase cascade activator may be administered as part of a pharmaceutical preparation containing suitable pharmaceutically acceptable carriers, comprising excipients and auxiliaries which facilitate processing of the compounds into preparations which can be used pharmaceutically. Preferably, the preparations, particularly those preparations which can be administered orally and which can be used for the preferred type of administration, such as tablets, dragees, and capsules, and also preparations which can be administered rectally,

such as suppositories, as well as suitable solutions for administration by injection or orally, contain from about 0.01 to 99 percent, preferably from about 0.25 to 75 percent of active compound(s), together with the excipient.

Also included within the scope of the present invention are the nontoxic pharmaceutically acceptable salts of the compounds of the present invention. Acid addition salts are formed by mixing a solution of the particular apoptosis inducer with a solution of a pharmaceutically acceptable non-toxic acid such as hydrochloric acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid, oxalic acid, and the like. Basic salts are formed by mixing a solution of the particular apoptosis inducers of the present invention with a solution of a pharmaceutically acceptable non-toxic base such as sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate, Tris, *N*-methyl-glucamine and the like.

The pharmaceutical compositions of the invention may be administered to any animal which may experience the beneficial effects of the compounds of the invention. Foremost among such animals are mammals, e.g., humans and veterinary animals, although the invention is not intended to be so limited.

The pharmaceutical compositions of the present invention may be administered by any means that achieve their intended purpose. For example, administration may be by parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, buccal, intrathecal, intracranial, intranasal or topical routes. Alternatively, or concurrently, administration may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

The pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, for example, by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus, pharmaceutical preparations for oral use can be obtained by combining the active compounds with solid excipients, optionally grinding the



resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

Suitable excipients are, in particular, fillers such as saccharides, for example lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or  
5 calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, as well as binders such as starch paste, using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents may be added such as the  
10 above-mentioned starches and also carboxymethyl-starch, crosslinked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable  
15 coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose  
20 preparations such as acetylcellulose phthalate or hydroxypropylmethyl-cellulose phthalate, are used. Dye stuffs or pigments may be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses. Other pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules  
25 made of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules can contain the active compounds in the form of granules which may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such  
30 as fatty oils, or liquid paraffin. In addition, stabilizers may be added. Possible

pharmaceutical preparations which can be used rectally include, for example, suppositories, which consist of a combination of one or more of the active compounds with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, or paraffin hydrocarbons. In addition, it is also possible to use gelatin rectal capsules which consist of a combination of the active compounds with a base. Possible base materials include, for example, liquid triglycerides, polyethylene glycols, or paraffin hydrocarbons. Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water soluble salts and alkaline solutions. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides or polyethylene glycol-400 (the compounds are soluble in PEG-400). Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, and include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers. In accordance with one aspect of the present invention, substituted 1,4-thiazepines or analogs thereof, or a pharmaceutically acceptable salt or prodrug thereof, are employed in topical and parenteral formulations and are used for the treatment of skin cancer.


The topical compositions of this invention are formulated preferably as oils, creams, lotions, ointments and the like by choice of appropriate carriers. Suitable carriers include vegetable or mineral oils; white petrolatum (white soft paraffin), branched chain fats or oils, animal fats and high molecular weight alcohol (greater than  $C_{12}$ ). The preferred carriers are those in which the active ingredient is soluble. Emulsifiers, stabilizers, humectants and antioxidants may also be included as well as agents imparting color or fragrance, if desired. Additionally, transdermal penetration enhancers can be employed in these topical formulations. Examples of such enhancers can be found in U.S. Patent Nos. 3,989,816 and 4,444,762.

-46-

Creams are preferably formulated from a mixture of mineral oil, self emulsifying beeswax and water in which mixture the active ingredient, dissolved in a small amount of an oil such as almond oil, is admixed. A typical example of such a cream is one which includes about 40 parts water, about 20 parts beeswax, about 40 parts mineral oil and about 1 part almond oil.

Ointments may be formulated by mixing a solution of the active ingredient in a vegetable oil such as almond oil with warm soft paraffin and allowing the mixture to cool. A typical example of such an ointment is one which includes about 30% almond oil and about 70% white soft paraffin by weight.

Accordingly, an aspect of the present invention is a method of treating a disorder responsive to the induction of apoptosis in an animal suffering from said disorder, which method comprises administering to the animal an effective amount of a compound of Formula I:



in which:

the dashed lines indicate optional unsaturation without violating valency rules;

$R^1$  is hydrogen,  $(C_{1-6})$ alkyl or  $-C(O)R^6$ , wherein  $R^6$  is as defined below, or is absent when a double bond exists between the nitrogen atom to which  $R^1$  is attached and an adjacent ring atom or  $R^1$  is as defined below;

$X^1$  is  $-NR^2$ -,  $-S$ -,  $-S(O)$ -,  $-S(O)_2$ - or  $-O$ -, wherein  $R^2$  is hydrogen or  $(C_{1-6})$ alkyl or is absent when a double bond exists between the nitrogen atom to which  $R^2$  is attached and an adjacent ring atom;

$A^1$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or


heterocycloalkyl each containing a total of 3 to 14 ring atoms, or A<sup>1</sup> together with R<sup>1</sup> and the atoms to which A<sup>1</sup> and R<sup>1</sup> are attached forms a fused polycyclic ring system selected from heteroaryl and unsaturated, partially unsaturated or saturated heterocycloalkyl in any case containing a total of 10 to 15 ring atoms, wherein A<sup>1</sup> may be substituted with a group selected from -R<sup>3</sup>, -X<sup>2</sup>OR<sup>3</sup>, -X<sup>2</sup>C(O)R<sup>3</sup>, -X<sup>2</sup>OC(O)R<sup>3</sup>, -X<sup>2</sup>C(O)OR<sup>3</sup>, -X<sup>2</sup>SR<sup>3</sup>, -X<sup>2</sup>S(O)R<sup>3</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>3</sup>, -X<sup>2</sup>NR<sup>3</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>3</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>3</sup>, -X<sup>2</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>3</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>3</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>3</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>3</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>3</sup> is -X<sup>2</sup>R<sup>5</sup> wherein X<sup>2</sup> is as defined above and R<sup>5</sup> is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, wherein each ring within A<sup>1</sup> and R<sup>5</sup> contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within A<sup>1</sup> and R<sup>5</sup> may be substituted further with 1 to 2 groups independently selected from (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo, with the provisos that only one of A<sup>1</sup> and R<sup>5</sup> is a fused polycyclic ring system;

A<sup>2</sup> is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein A<sup>2</sup> may be substituted with a group selected from -R<sup>8</sup>, -X<sup>2</sup>OR<sup>8</sup>, -X<sup>2</sup>C(O)R<sup>8</sup>, -X<sup>2</sup>OC(O)R<sup>8</sup>, -X<sup>2</sup>C(O)OR<sup>8</sup>, -X<sup>2</sup>SR<sup>8</sup>, -X<sup>2</sup>S(O)R<sup>8</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>8</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>8</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>8</sup>,

wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$  is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^2$  and  $R^8$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said heterocycloalkylene, carbocycloalkyl and heterocycloalkyl rings within  $A^2$  and  $R^8$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^2$  and  $R^8$  is a fused polycyclic ring system; and

$A^3$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-R^9$ ,  $-X^2OR^9$ ,  $-X^2C(O)R^9$ ,  $-X^2OC(O)R^9$ ,  $-X^2C(O)OR^9$ ,  $-X^2SR^9$ ,  $-X^2S(O)R^9$ ,  $-X^2S(O)_2R^9$ ,  $-X^2NR^4R^9$ ,  $-X^2NR^4C(O)R^9$ ,  $-X^2NR^4C(O)OR^9$ ,  $-X^2C(O)NR^4R^9$ ,  $-X^2NR^4C(O)NR^4R^9$ ,  $-X^2NR^4C(NR^4)NR^4R^9$ ,  $-X^2NR^4S(O)_2R^9$  and  $-X^2S(O)_2NR^4R^9$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^9$  is  $-X^2R^{10}$  wherein  $X^2$  is as defined above and  $R^{10}$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within

$A^3$  and  $R^{10}$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within  $A^3$  and  $R^{10}$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^3$  and  $R^{10}$  is a fused polycyclic ring system; or an *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers, or a pharmaceutically acceptable salt thereof; with the proviso that when said compound is of Formula II(a):



II(a)

then  $A^3$  is other than:

- (a) benzo[1,3]dioxolyl;
- (b) phenyl which is mono-substituted by bromo, hydroxy, methyl or isopropyl; and
- (c) phenyl which is substituted by at least one of chloro and methoxy and not substituted by methylsulfanyl, amino, methylamino or dimethylamino; preferably wherein said disorder is an autoimmune disease, in particular rheumatoid arthritis, or inflammation or inflammatory bowel disease, in particular, wherein said disorder is psoriasis or a skin disease.

Another aspect of the present invention is a method for treating or preventing cancer, comprising administering to an animal in need of such treatment an effective amount of a compound of Formula I-as defined

immediately above; particularly wherein said cancer is selected from the group consisting of Hodgkin's disease, non-Hodgkin's lymphoma, acute and chronic lymphocytic leukemias, multiple myeloma, neuroblastoma, breast carcinoma, ovarian carcinoma, lung carcinoma, Wilms' tumor, cervical carcinoma, testicular carcinoma, soft-tissue sarcoma, chronic lymphocytic leukemia, primary macroglobulinemia, bladder carcinoma, chronic granulocytic leukemia, primary brain carcinoma, malignant melanoma, small-cell lung carcinoma, stomach carcinoma, colon carcinoma, malignant pancreatic insulinoma, malignant carcinoid carcinoma, choriocarcinoma, mycosis fungoides, head and neck carcinoma, osteogenic sarcoma, pancreatic carcinoma, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinoma, thyroid carcinoma, esophageal carcinoma, malignant hypercalcemia, cervical hyperplasia, renal cell carcinoma, endometrial carcinoma, polycythemia vera, essential thrombocytosis, adrenal cortex carcinoma, skin cancer and prostatic carcinoma.


Another aspect of the present invention is a method for the treatment of drug resistant cancer, comprising administering to an animal in need of such treatment an effective amount of a compound of compound of Formula I as defined immediately above.

Preferred means for practicing any of the above methods comprises further administering to said animal at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent; preferably wherein said known cancer therapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin, imitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin®, Rituxan® and alanosine.

Preferred means for practicing any of the above methods comprises further treating said animal with radiation-therapy.

Preferred means for practicing any of the above methods comprises administering the compound of Formula I after surgical treatment for cancer.

5 Preferred means for practicing any of the above methods comprises administering a compound of Formula I which when said compound is selected from the group consisting of Formula II(a) and II(b):



then A<sup>3</sup> is other than:

- 10 (a) benzo[1,3]dioxolyl;
- (b) phenyl which is mono-substituted by bromo, nitro, hydroxy, methyl or isopropyl; and
- (c) phenyl which is substituted by at least one of Cl and methoxy and not substituted by methylsulfanyl, amino, methylamino and dimethylamino.


15 A further preferred means for practicing any of the above methods comprises administering a compound of Formula I which when said compound is selected from the group consisting of Formula II(a) and II(b), then A<sup>3</sup> is other than benzo[1,3]dioxol-5-yl, 2,3-dihydro-benzo[1,4]dioxinyl or phenyl which is substituted by at least one of bromo, chloro, hydroxy, nitro, methoxy and

20 (C<sub>1-3</sub>)alkyl


A further preferred means for practicing any of the above methods comprises administering a compound of Formula I in which A<sup>1</sup> of said compound is a group selected from Formulae (a), (b), (c), (d) and (e):




-52-




(a)




(b)



(c)



(d)




(e)

in which  $R^7$  is hydrogen or methyl,  $R^{11}$  is hydrogen or  $(C_{1-6})$ alkyl and the free valance is attached to  $A^2$ , or

$A^2$  and  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a group of Formula (g):

5




(g)


wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ ; and

$A^2$  of said compound is as defined above or is a group selected from Formulae (h), (i), (j), (k), (l) and (m):


-53-




(h)




(i)




(j)



(k)



(l)



(m)

in which  $n$  is 1 or 2 and  $R^1$  is acetyl or trifluoroacetyl.

A further preferred means for practicing any of the above methods comprises administering a compound of Formula I in which  $A^3$  is phenyl or heteroaryl containing a total of 5 to 9 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-R^9$ ,  $-X^2OR^9$ ,  $-X^2SR^9$  and  $-X^2S(O)_2R^9$ , wherein  $R^9$  is  $-X^2R^{10}$ ,  $X^2$  is a bond or  $(C_{1-6})$ alkylene and  $R^{10}$  is phenyl or heteroaryl containing a total of 5 to 6 ring atoms, wherein each ring within  $A^3$  and  $R^{10}$  may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, halo, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)_2R^6$  and  $-X^2NR^4R^4$ , wherein  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl.

A further preferred means for practicing any of the above methods comprises administering a compound of Formula I, wherein said compound is selected from the group consisting of:

4-hydroxy-3-[7-(2-methoxy-4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

2-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

4-hydroxy-3-[7-(4-methanesulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

10 2-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

3-hydroxy-2-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-cyclohex-2-enone;

3-hydroxy-2-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

15 4-hydroxy-6-methyl-3-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-5,6-dihydro-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-5,6-dihydro-pyran-2-one; and

20 3-[7-(2,4-dimethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

A further preferred means for practicing any of the above methods comprises administering a compound of Formula I, wherein said compound is selected from the group consisting of:

25 3-[4-acetyl-7-(2,4-dimethoxy-phenyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,4-dimethoxy-phenyl)-4-(2,2,2-trifluoro-ethanoyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

1-[7-(2,4-dimethoxy-phenyl)-5-(3-fluoro-4-methoxyphenyl)-[1,4]thiazepan-4-yl]-ethanone;

4-hydroxy-6-methyl-3-[7-(3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(5-ethyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 3-[7-(1-benzyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3[7-(2-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

10 4-hydroxy-6-methyl-3[7-(3-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3[7-(4-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-[3-(3-trifluoromethyl-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

15 3-[7-[3-(3,4-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

20 4-hydroxy-6-methyl-3-{7-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

3-{7-[5-(2-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[5-(3-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

25 3-{7-[5-(4-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-{7-[5-(chloro-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

30 3-[7-(4-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(1-benzenesulfonyl-1*H*-pyrrol-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 4-hydroxy-6-methyl-3-[7-(3-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

10 4-hydroxy-6-methyl-3-[7-(1-methyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(chloro-methyl-trifluoromethyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[1-(2,4-difluoro-benzenesulfonyl)-1*H*-pyrrol-2-yl]}-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

15 3-(7-[2,2']bithienyl-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]}-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

20 3-{7-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]}-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-chloro-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

25 3-[7-(4,5-dibromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2-chloro-5-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

30 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methylsulfanyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(5-chloro-1-methyl-3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 3-[4-(2,4-dimethoxy-phenyl)-4,5-dihydro-3*H*-benzo[*b*][1,4]diazepin-2-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 3-[7-2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-1*H*-quinolin-2-one;

4-hydroxy-6-methyl-3-[7-(4-trifluoromethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(bis-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

15 3-[7-(4-dimethylamino-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-hydroxy-2-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

20 3-[7-(2,4-dimethoxy-phenyl)-1-oxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>4</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methoxy-pyran-2-one;

10-(2,4-dimethoxy-phenyl)-3-methyl-7,8-dihydro-10*H*-2,5-dioxa-9-thia-6a-aza-cyclohepta[*a*]naphthalene-1,6-dione;

3-[7-(2,4-dimethoxy-phenyl)-1,1-dioxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>6</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

25 3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl-2,3-dihydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one;

30 2-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone; and

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one.

A further preferred means for practicing any of the above methods comprises administering a compound of Formula I, wherein said compound is selected from the group consisting of:

4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(4-ethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(3-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[7-(2-bromo-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,3-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

6-methyl-3-(2-*p*-tolyl-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl)-pyran-2-one;

4-hydroxy-6-methyl-3-[2-(4-methylsulfanyl-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-pyran-2-one.

A further preferred means for practicing any of the above methods comprises administering a compound of Formula I, wherein said compound is selected from the group consisting of:

4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-3-[7-(4-chloro-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methylpyran-2-one; and


5 4-hydroxy-3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one.

### Chemistry:

#### Processes for Making Compounds of the Invention:

10 Compounds of the invention may be prepared by proceeding as in Scheme 1.

Scheme 1



15 in which the dashed line indicates optional unsaturation, X<sup>4</sup> is -NR<sup>2</sup> or SH, R<sup>12</sup> is -NHR<sup>2</sup> or -SH, R<sup>13</sup> and R<sup>14</sup> independently are hydrogen or any of the optional substituents defined for A<sup>2</sup> in the Summary of the Invention or R<sup>13</sup> and R<sup>14</sup> together with the atoms to which R<sup>13</sup> and R<sup>14</sup> are attached form a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 9 ring



atoms, heteroaryl containing a total of 5 to 9 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 9 ring atoms and the compound of Formula 4 may be optionally substituted.

5           Compounds of the invention may be prepared by reacting a compound of Formula 3 with a compound of Formula 4. The reaction is carried out in a suitable solvent (e.g., ethanol) at between 60 and 80°C and requires 10 minutes to 20 hours to complete. A detailed description for the preparation of a compound of Formula I by the methods described above is set forth in  
10           Example 2, *infra*.

          Compounds of Formula 3 can be prepared by reacting a ketone of Formula I with an aldehyde of Formula 2. The reaction is carried out in an appropriate solvent (e.g., ethanol) and in the presence of a suitable base (e.g., piperidine) at between 80 to 85°C and requires 8 to 24 hours to complete. A  
15           detailed description for the preparation of a compound of Formula I by the methods described above is set forth in Reference 5.1, *infra*.

          Compounds of Formula I in which A<sup>2</sup> is [1,4]thiazepanylene wherein R<sup>1</sup> is -C(O)R<sup>6</sup> can be prepared by reducing a compound of Formula I in which A<sup>2</sup> is 2,3,6,7-tetrahydro-[1,4]thiazepinylene to the corresponding thiazepane and then  
20           condensing the thiazepane with an anhydride of the Formula O[C(O)R<sup>6</sup>]<sub>2</sub>. The reduction is carried out in a suitable solvent (e.g., ethanol) and in the presence of a suitable reducing agent (e.g., sodium borohydride) at 35 and 50°C and requires 0.5 to 6 hours to complete. The condensation reaction is carried out in the presence of suitable base (e.g., diisopropylethylamine (DIPEA)) and requires 0.5  
25           to 14 hours to complete. A detailed description for the preparation of a compound of Formula I by the methods described above is set forth in Example 1, *infra*.

          Compounds of Formula I in which A<sup>2</sup> is 1-oxo-2,3,6,7-tetrahydro-1H-1λ<sup>4</sup>-[1,4]thiazepinylene can be prepared by oxidizing a compound of Formula I in  
30           which A<sup>2</sup> is 2,3,6,7-tetrahydro-[1,4]thiazepinylene. The oxidation is carried out

in a suitable solvent (e.g., acetic acid) and in the presence of a suitable oxidizing agent (e.g., hydrogen peroxide) at ambient temperature and requires 2 to 3 hours to complete. Compounds of Formula I in which A<sup>2</sup> is 1,1-dioxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>4</sup>-[1,4]thiazepinylenes can be prepared by proceeding as described above, but heating the reaction at approximately 70°C for 2 to 3 hours. Detailed descriptions for the preparation of a compound of Formula I by the methods described above are set forth in Examples 3 and 5, *infra*.

Compounds of Formula I in which A<sup>2</sup> is [1,4]thiazepanylene wherein R<sup>1</sup> and A<sup>1</sup> and the atoms to which A<sup>1</sup> and R<sup>1</sup> are attached together with A<sup>2</sup> form a group of Formula (h) can be prepared by reacting a corresponding compound of Formula I in which A<sup>2</sup> is 2,3,6,7-tetrahydro-[1,4]thiazepinylenes and A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl with phosgene. The reaction is carried out in a suitable solvent (e.g., ethylene dichloride) and in the presence of a suitable acylation catalyst (e.g., 4-(dimethylamino)pyridine) and in the presence of suitable base (e.g., DIPEA) at ambient temperature and requires 0.5 to 1 hours to complete. A detailed description for the preparation of a compound of Formula I by the methods described above is set forth in Example 4, *infra*.

Compounds of Formula I in which A<sup>2</sup> is 2,3-dihydro-[1,4]thiazepinylenes can be prepared by dehydrogenation of a corresponding compound of Formula I in which A<sup>2</sup> is 2,3,6,7-tetrahydro-[1,4]thiazepinylenes. The reaction is carried out in an appropriate solvent (e.g., toluene) and in the presence of a suitable catalyst (e.g., 2,3-dichloro-5,6-dicyano-1,4-benzoquinone) at 60 to 80°C and requires 20 to 30 minutes to complete. A detailed description for the preparation of a compound of Formula I by the methods described above is set forth in Example 5, *infra*.

## EXPERIMENTALS

### REFERENCE 1

#### 3-Acetyl-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one

4-Hydroxy-6-methyl-5,6-dihydro-pyran-2-one (2 g, 15.6 mmol) was combined with methylene chloride (50 mL) in a 250 mL flask and then acetic acid (17.5 M, 2 eq, 31 mmol, 1.77 mL) was added to the mixture. The mixture was cooled in an ice bath and then dicyclohexylcarbodiimide (1.3 eq, 20.3 mmol, 4.19 g) was added portionwise, followed by the addition of 4-(dimethylamino)pyridine (0.05 eq, 0.78 mmol, 88 mg) to the mixture. A sufficient amount of methylene chloride was added to ensure easy stirring and the reaction was monitored by TLC (silica gel, hexane-ethyl acetate-methylene chloride-acetone (3:3:3:1 v/v)), HPLC and LCMS as the intermediate acetic acid 2-methyl-6-oxo-3,6-dihydro-2H-pyran-4-yl ester was formed. The mixture was stirred overnight at room temperature and then toluene (20 mL) was added. The mixture was heated to 60°C and after 48 hours the mixture was filtered, concentrated and purified by flash column chromatography to provide 3-acetyl-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one (1.02 g, 38%) as a white solid, mp 96–98°C (lit., mp 97–98°C). LCMS: MH<sup>+</sup> 171.0.

### REFERENCE 2

#### 2-Acetyl-3-hydroxy-cyclohex-2-enone

3-Hydroxy-cyclohex-2-enone (1.08 g, 9.64 mmol) was combined with methylene chloride (10 mL) in a 250 mL flask and acetic acid (1.4 eq, 13.75 mmol, 0.78 mL) was added to the mixture. The mixture was cooled in an ice bath and dicyclohexylcarbodiimide (1.3 eq, 12 mmol, 2.5 g) was added portionwise, followed by the addition of 4-(dimethylamino)pyridine (0.05 eq, 0.45 mmol, 50 mg). A sufficient amount of methylene chloride was added to

ensure easy stirring and the reaction was monitored by TLC (silica gel, hexane-ethyl acetate-methylene chloride- acetone (3:3:3:1 v/v)), HPLC and LCMS and the intermediate acetic acid 3-oxo-cyclohex-1-enyl ester was formed. The mixture was stirred overnight at room temperature and then toluene (20 mL) was added. The mixture heated to 60°C and after 72 hours the mixture was filtered, concentrated and purified by flash column chromatography to provide 2-acetyl-3-hydroxy-cyclohex-2-enone (1.19 g, 80%) as a colorless liquid. LCMS: MH<sup>+</sup> 154.8.

## REFERENCE 3

3-Acetyl-4-hydroxy-1*H*-quinolin-2-one

A solution of 2-amino-benzoic acid methyl ester (1.51 g, 10 mmol) and triethylamine (97.2 mmol/mL, 0.0194 mL, 0.14 mmol, 0.014 eq) in toluene (4 mL) was heated to 60°C and then a solution of 4-methylene-oxetan-2-one (0.84 g, 10 mmol) in toluene (2 mL) was added to the solution over 15 minutes. The reaction was heated at 80°C for 6 hours and then at 50°C for 16 hours. Progress of the reaction was monitored by TLC (silica gel, hexane-ethyl acetate (7:3 v/v)), analytical HPLC and LCMS and upon its completion the mixture was partitioned between ethyl acetate and aqueous hydrochloric acid (1 N). The organic layer was washed with water, saturated sodium bicarbonate, water and then brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give an orange solid. The solid was crystallized from methylene chloride /ethyl acetate and hexane to provide 2-(3-oxo-butanoylamino)-benzoic acid methyl ester (1.83 g, 78%) as large colorless prisms, mp 81-83 °C (lit., mp 79-80°C). LCMS MH<sup>+</sup> 235.6.

2-(3-Oxo-butanoylamino)-benzoic acid methyl ester (383 mg, 1.63 mmol) was combined with diethyl ether (10 mL) and methanol (5 mL) and the mixture was stirred rapidly while a solution of sodium methoxide (25 % solution, 1.63 mmol, 0.45 mL) in methanol (3 mL) was added over 10 minutes. The reaction then was heated at 40°C overnight. Progress of the reaction was monitored by

TLC (silica gel, hexane : EtOAc : CH<sub>2</sub>Cl<sub>2</sub> : acetone (3:3:3:1)) and analytical HPLC and upon its completion the mixture was acidified with 1 N sulfuric acid to form a solid. The solid was collected by filtration and then crystallized from hot acetic acid / acetonitrile / water to provide 3-acetyl-4-hydroxy-1*H*-quinolin-2-one (250 mg, 76%) as small colorless prisms, mp 258–262°C (dec) (lit., 259°C). LCMS MH<sup>+</sup> 204.2.

## REFERENCE 4

## 2-Methoxy-4-methylsulfanyl-benzaldehyde

2-Methoxy-4-methylsulfanyl-benzoic acid (2 g, 10.09 mmol) was dissolved in dry THF (20 mL) and the solution was stirred while heated to 60°C under nitrogen and then borane-methylsulfide complex (1.7 eq, 1.7 mL, 17.5 mmol) was added very slowly dropwise via a syringe. The progress of the reaction was followed by both TLC and analytical HPLC and when complete (3 hours) the mixture was allowed to cool to room temperature, diluted with water (10 mL) added extremely slowly dropwise under nitrogen. Potassium carbonate (1 g) was added and after stirring the mixture for 30 minutes ethyl acetate (50 ml) was added. The organic layer was separated, washed with water, 2 N hydrochloric acid, water and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and then concentrated to a near colorless oil. The residue was triturated with hexane and product was purified from the resulting crystals by flash column chromatography to provide (2-methoxy-4-methylsulfanyl-phenyl)-methanol (934 mg, 51%) as colorless crystals. LCMS: M<sup>-</sup> 182.8.

Pyridine (4.8 mL, 60 mmol, 12 eq) was added under nitrogen to a mixture of chromium trioxide (fresh, 2 g, 20 mmol) in dry methylene chloride (15 mL) cooled over an ice bath. The mixture was stirred at 0°C for 1 hour and Celite powder (1 g) and a solution of (2-methoxy-4-methylsulfanyl-phenyl)-methanol (934 mg, 5.07 mmol) in methylene chloride (10 mL) was added. The progress of the reaction was followed by TLC (silica gel, hexane-EtOAc-CH<sub>2</sub>Cl<sub>2</sub>-acetone

(3:3:3:1 v/v) and hexane-ethyl acetate (7:3 v/v)) and analytical HPLC and when complete, the mixture was applied to a silica gel column (made up in hexane) and the column eluted with methylene chloride. Pure fractions were combined and concentrated to provide 2-methoxy-4-methylsulfanyl-benzaldehyde (730 mg, 79%). LCMS:  $MH^+$  183.2.

## REFERENCE 5

3-[7-(2,4-Dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-  
4-hydroxy-6-methyl-pyran-2-one

### Step 5.1

3-Acetyl-4-hydroxy-6-methyl-pyran-2-one (5.07 g, 30.15 mmol) and 2,4-dimethoxybenzaldehyde (5.03 g, 30.27 mmol) were combined in a 250 mL round bottom flask. Absolute ethanol (20 mL) and piperidine (0.10 eq, 0.30 mL, 3.0 mmol) were added and the mixture heated to between 80 and 85 °C. The reaction was monitored by TLC (silica gel, hexane-ethyl acetate (1:1 v/v) and methylene chloride-ethyl acetate-acetone (5:5:1 v/v)) and analytical HPLC. After 20 hours (>95% completion) the reaction mixture was cooled to room temperature to provide 3-[3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one as a bright orange crystalline precipitate. MS  $M^+$  317. NMR ( $CDCl_3$ -TMS):  $\delta$  8.31 (m, 2H), 7.69 (d, 1H,  $J=9$  Hz), 6.54 (d, 1H,  $J=9$  Hz), 6.45 (s, 1H), 5.93 (s, 1H), 3.91 (s, 3H), 3.87 (s, 3H), 2.26 (s, 3H) NMR ( $DMSO-d_6$ ):  $\delta$  8.11 (s, 2H), 7.62 (d, 1H,  $J=9$  Hz), 6.63 (bs, 2H), 6.23 (s, 1H), 3.87 (s, 3H), 3.82 (s, 3H), 2.28 (s, 3H).

### Step 5.2

The crude reaction mixture prepared in Step 5.1 was diluted with absolute ethanol (5 mL). 2-Aminoethanethiol (2.31 g, 30 mmol fresh dry material) was added and the mixture heated to 75°C. The reaction was monitored by TLC (silica gel, hexane-ethyl acetate (1:1 v/v) and methylene chloride-ethyl acetate-

acetone (5:5:1 v/v)) and analytical HPLC. After 1.5 hours (>90% completion) the reaction was allowed to cool to room temperature to form a yellow crystalline solid. The solid material was collected by filtration and washed with ethanol-diethyl ether (1:1 v/v) and then hexane. Product was purified by column chromatography and then crystallization (methylene chloride-ethanol) to provide 3-[7-(2,4-dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (6.81 g, 61%). LCMS:  $MH^+$  375.6. Elemental Analysis: Calc: C 60.78, H 5.64, N 3.73 Found: 60.73, H 5.66, N 3.73.  $^1H$  NMR ( $CDCl_3$ -TMS):  $\delta$  14.3 (bs, 1H), 7.23 (m, 1H), 6.46 (m, 2H), 5.69 (s, 1H), 4.76 (d, 1H,  $J=13$  Hz), 4.47 (d, 1H,  $J=11$  Hz), 4.17 (m, 1H), 4.00 (m, 1H), 3.83 (s, 3H), 3.79 (s, 3H), 3.52 (t, 1H,  $J=11.5$  Hz), 3.00 (t, 1H,  $J=11.5$  Hz), 2.78 (dd, 1H,  $J=11.5, 5$  Hz), 2.09 (s, 3H).  $^1H$  NMR ( $DMSO-d_6$ ):  $\delta$  13.7 (bs, 1H), 7.23 (dd, 1H,  $J=9, 3$  Hz), 6.5 (m, 2H), 5.7 (s, 1H), 4.37 (d, 1H,  $J=13$  Hz), 4.28 (d, 1H,  $J=10$  Hz), 4.08 (m, 2H), 3.73 (s, 6H), 3.61 (t, 1H,  $J=11$  Hz), 2.79 (m, 2H), 2.03 (s, 3H).  $^{13}C$  NMR JEOL, ( $DMSO-d_6$ ):  $\delta$  183.84, 178.36, 163.42, 162.89, 160.32, 157.33, 128.28, 123.29, 107.76, 105.46, 99.16, 95.78, 56.26, 55.77, 46.0, 38.90, 34.19, 29.07, 19.68.

Proceeding as in Reference 5, Step 5.1, but substituting 3-acetyl-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 3-[(E)-3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 2-acetyl-3-hydroxy-cyclohex-2-enone for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 2-[3-(2,4-dimethoxy-phenyl)-acryloyl]-3-hydroxy-cyclohex-2-enone.

Proceeding as in Reference 5, Step 5.1, but substituting 3-acetyl-4-hydroxy-1H-quinolin-2-one for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 3-[(E)-3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-1H-quinolin-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 2-methoxy-4-methylsulfanyl-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-

hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-phenyl-1*H*-pyrazole-4-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(3-phenyl-1*H*-pyrazol-4-yl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-acetyl-4-methoxy-6-methyl-pyran-2-one for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 3-[(E)-3-(2,4-dimethoxy-phenyl)-acryloyl]-4-methoxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-ethyl-thiene-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(5-ethyl-thien-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 1-benzyl-1*H*-indole-3-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(1-benzyl-1*H*-indol-3-yl)acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 2-trifluoromethylsulfanyl-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(2-trifluoromethylsulfanyl-phenyl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-trifluoromethylsulfanyl-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(3-trifluoromethylsulfanyl-phenyl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 4-trifluoromethylsulfanyl-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(4-trifluoromethylsulfanyl-phenyl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-(3-trifluoromethyl-phenoxy)-benzaldehyde for 2,4-dimethoxybenzaldehyde,



provided 4-hydroxy-6-methyl-3-[(E)-3-[3-(3-trifluoromethyl-phenoxy)-phenyl]-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-(3,4-dichloro-phenoxy)-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-[3-(3,4-dichloro-phenoxy)-phenyl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-(3,5-dichloro-phenoxy)-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-[3-(3,5-dichloro-phenoxy)-phenyl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-(3-trifluoromethyl-phenyl)-furan-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-(2-chloro-phenyl)-furan-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-[5-(2-chloro-phenyl)-furan-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but 5-(3-chloro-phenyl)-furan-2-carbaldehyde substituting for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-[5-(3-chloro-phenyl)-furan-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-(4-chloro-phenyl)-furan-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-[5-(4-chloro-phenyl)-furan-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-(chloro-trifluoromethyl-phenyl)-furan-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-[5-(chloro-trifluoromethyl-phenyl)-furan-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 4-bromo-thiene-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(4-bromo-thien-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-bromo-thiene-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(5-bromo-thien-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but 1-benzenesulfonyl-1*H*-pyrrole-2-carbaldehyde substituting for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(1-benzenesulfonyl-1*H*-pyrrol-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-methyl-thiene-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(3-methyl-thien-2-yl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-methyl-thiene-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(5-methyl-thien-2-yl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 1-methyl-1*H*-indole-3-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(1-methyl-1*H*-indol-3-yl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting chloro-methyl-trifluoromethyl-1*H*-pyrazole-4-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(chloro-methyl-trifluoromethyl-1*H*-pyrazol-4-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 1-(2,4-difluorobenzenesulfonyl)-1*H*-pyrrole-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-[1-(2,4-difluorobenzenesulfonyl)-1*H*-pyrrol-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting [2,2']bithienyl-5-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-[2,2']bithienyl-5-yl-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 1-(3,5-dichlorophenyl)-1*H*-pyrrole-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-

{(E)-3-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]-acryloyl}-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 1-(4-chloro-phenyl)-1*H*-pyrrole-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-  
5 {(E)-3-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]-acryloyl}-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-chloro-1*H*-indole-3-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(5-chloro-1*H*-indol-3-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazole-5-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided  
10 4-hydroxy-6-methyl-3-[(E)-3-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 4,5-dibromothi-  
15 ene-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(4,5-dibromo-thien-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting chloro-trifluoromethyl-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(chloro-trifluoromethyl-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-acetyl-4-  
20 hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 3-[(E)-3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-methylsulfanyl-  
25 thiene-2-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(5-methylsulfanyl-thien-2-yl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 5-chloro-1-methyl-3-phenyl-1*H*-pyrazole-4-carbaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(5-chloro-1-methyl-3-phenyl-1*H*-pyrazol-4-yl)-acryloyl]-4-  
30 hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 2-acetyl-3-hydroxy-cyclohex-2-enone for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 2-[3-(2,4-dimethoxy-phenyl)-acryloyl]-3-hydroxy-cyclohex-2-enone.

Proceeding as in Reference 5, Step 5.1, but substituting 1-(3-fluoro-4-methoxy-phenyl)-ethanone for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided (E)-1-(3-fluoro-4-methoxy-phenyl)-3-(2-hydroxy-4-methoxy-phenyl)-propenone.

Proceeding as in Reference 5, Step 5.1, but substituting 4-dimethylamino-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(4-dimethylamino-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 3-acetyl-4-hydroxy-1*H*-quinolin-2-one for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 3-[(E)-3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-1*H*-quinolin-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 4-trifluoromethoxy-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-6-methyl-3-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting bis-trifluoromethyl-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(bis-trifluoromethyl-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 4-dimethylamino-2-methoxy-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(4-dimethylamino-2-methoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 4-methanesulfonyl-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-3-[(E)-3-(4-methanesulfonyl-phenyl)-acryloyl]-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 2,4-diethoxy-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 3-[(E)-3-(2,4-diethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 2,4-diethoxybenzaldehyde for 2,4-dimethoxybenzaldehyde and 3-acetyl-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 3-[(E)-3-(2,4-diethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 4-dimethylamino-benzaldehyde for 2,4-dimethoxybenzaldehyde and 3-acetyl-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 3-[(E)-3-(4-dimethylamino-phenyl)-acryloyl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 2,4-diethoxybenzaldehyde for 2,4-dimethoxybenzaldehyde and 2-acetyl-3-hydroxy-cyclohex-2-enone for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 2-[(E)-3-(2,4-diethoxy-phenyl)-acryloyl]-3-hydroxy-cyclohex-2-enone.

Proceeding as in Reference 5, Step 5.1, but substituting 2,4-diethoxybenzaldehyde for 2,4-dimethoxybenzaldehyde and 2-acetyl-3-hydroxy-cyclohex-2-enone for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 2-[(E)-3-(2,4-diethoxy-phenyl)-acryloyl]-3-hydroxy-cyclohex-2-enone.

Proceeding as in Reference 5, Step 5.1, but substituting 2,3,4-trimethoxybenzaldehyde for 2,4-dimethoxybenzaldehyde and 2-acetyl-3-hydroxy-cyclohex-2-enone for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided 3-hydroxy-2-[(E)-3-(2,3,4-trimethoxy-phenyl)-acryloyl]-cyclohex-2-enone.


Proceeding as in Reference 5, Step 5.1, but substituting 2-methoxy-4-methylsulfanyl-benzaldehyde for 2,4-dimethoxybenzaldehyde, provided 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one.

Proceeding as in Reference 5, Step 5.1, but substituting 2,3,4-trimethoxybenzaldehyde for 2,4-dimethoxybenzaldehyde and 3-acetyl-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 3-acetyl-4-hydroxy-6-methyl-pyran-2-one, provided

4-hydroxy-6-methyl-3-[(E)-3-(2,3,4-trimethoxy-phenyl)-acryloyl]-5,6-dihydro-pyran-2-one.

### EXAMPLE 1

3-[4-Acetyl-7-(2,4-dimethoxy-phenyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one  
(Compound 1)




3-[7-(2,4-Dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (187 mg, 0.5 mmol), prepared as in Reference 5, was combined with absolute ethanol (5 mL) and sodium borohydride (20 mg, 0.526 mmol) and the mixture was warmed on a hot plate to form a solution. The progress of the reaction was followed by analytical HPLC and LCMS and when complete (5 hours) acetic anhydride (1.5 mL, 30 eq) and DIPEA (0.6 mmol, 0.1 mL) were added. The reaction was stirred at room temperature overnight. Product was isolated by preparative HPLC (RPC<sub>18</sub> column, acetonitrile/water containing 0.1% HCl) to provide 3-[4-acetyl-7-(2,4-dimethoxy-phenyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one (25 mg). LCMS: MH<sup>+</sup> 420.2.

Proceeding as in Example 1, but substituting trifluoroacetic anhydride for acetic anhydride, provided 3-[7-(2,4-dimethoxy-phenyl)-4-(2,2,2-trifluoroethanoyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 2); LCMS: MH<sup>+</sup> 474.0.

Proceeding as in Example 1, but substituting 7-(2,4-dimethoxy-phenyl)-5-(3-fluoro-4-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepine for 3-[7-(2,4-dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one, provided 1-[7-(2,4-dimethoxy-phenyl)-5-(3-fluoro-4-methoxy-phenyl)-[1,4]thiazepan-4-yl]-ethanone (Compound 3); LCMS:  $MH^+$  420.2.

## EXAMPLE 2

4-Hydroxy-3-[7-(2-methoxy-4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one  
(Compound 4)



4-Hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one (41.8 mg, 0.126 mmol), prepared as in Reference 5, Step 5.1, was dissolved in absolute ethanol (5 mL) and then 2-aminoethanethiol (9.8 mg, 0.126 mmol) was added to the solution. The mixture was stirred at 80°C. The progress of the reaction was followed by analytical HPLC and when complete (approximately 12 hours) the solvent was removed *in vacuo*. Product was purified by silica flash column chromatography to provide 4-hydroxy-3-[7-(2-methoxy-4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one (20.5 mg) as a pale, yellow solid. LCMS:  $MH^+$  392.6.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(3-phenyl-1H-pyrazol-4-yl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(3-phenyl-1H-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 5); LCMS:  $MH^+$  383.

Proceeding as in Example 2, but substituting 3-[(E)-3-(2,4-dimethoxy-phenyl)-acryloyl]-4-methoxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-methoxy-6-methyl-pyran-2-one (Compound 6); MS  $MH^+$  390.

Proceeding as in Example 2, but substituting 3-[(E)-3-(5-ethyl-thien-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(5-ethyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 7); LCMS:  $MH^+$  349.9.

Proceeding as in Example 2, but substituting 3-[(E)-3-(1-benzyl-1*H*-indol-3-yl)acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(1-benzyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 8); LCMS:  $MH^+$  445.2.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(2-trifluoromethylsulfanyl-phenyl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(2-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 9); LCMS:  $MH^+$  416.4.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(3-trifluoromethylsulfanyl-phenyl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(3-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 10); LCMS:  $MH^+$  416.6.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(4-trifluoromethylsulfanyl-phenyl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(4-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 11); LCMS:  $MH^+$  416.5.



Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-[3-(3-trifluoromethyl-phenoxy)-phenyl]-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-{7-[3-(3-trifluoromethyl-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one (Compound 12); LCMS: MH<sup>+</sup> 476.2.

Proceeding as in Example 2, but substituting 3-[(E)-3-[3-(3,4-dichloro-phenoxy)-phenyl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-{7-[3-(3,4-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one (Compound 13); LCMS: MH<sup>+</sup> 475.8.

Proceeding as in Example 2, but substituting 3-[(E)-3-[3-(3,5-dichloro-phenoxy)-phenyl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-{7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one (Compound 14); LCMS: MH<sup>+</sup> 475.9.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-{7-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one (Compound 15); LCMS: MH<sup>+</sup> 450.1.

Proceeding as in Example 2, but substituting 3-[(E)-3-[5-(2-chloro-phenyl)-furan-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-{7-[5-(2-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one (Compound 16); LCMS: MH<sup>+</sup> 416.1.

Proceeding as in Example 2, but substituting 3-[(E)-3-[5-(3-chloro-phenyl)-furan-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-[5-(3-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 17); LCMS: MH<sup>+</sup> 416.1.

Proceeding as in Example 2, but substituting 3-[(E)-3-[5-(4-chloro-phenyl)-furan-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-[5-(4-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 18); LCMS: MH<sup>+</sup> 416.1.

Proceeding as in Example 2, but substituting 3-[(E)-3-[5-(2-chloro-5-trifluoromethyl-phenyl)-furan-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-[5-(2-chloro-5-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 19); LCMS: MH<sup>+</sup> 483.9.

Proceeding as in Example 2, but substituting 3-[(E)-3-(4-bromo-thien-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(4-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 20); LCMS: MH<sup>+</sup> 401.8.

Proceeding as in Example 2, but substituting 3-[(E)-3-(5-bromo-thien-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(5-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 21); LCMS: MH<sup>+</sup> 401.8.

Proceeding as in Example 2, but substituting 3-[(E)-3-(1-benzenesulfonyl-1H-pyrrol-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(1-benzenesulfonyl-1H-pyrrol-2-yl)-2,3,6,7-tetrahydro-

[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 22); LCMS:  $MH^+$  445.5.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(3-methyl-thien-2-yl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(3-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 23); LCMS:  $MH^+$  336.0.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(5-methyl-thien-2-yl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(5-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 24); LCMS:  $MH^+$  336.0.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(1-methyl-1*H*-indol-3-yl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(1-methyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 25); LCMS:  $MH^+$  369.2.

Proceeding as in Example 2, but substituting 3-[(E)-3-(chloro-methyl-trifluoromethyl-1*H*-pyrazol-4-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(chloro-methyl-trifluoromethyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 26); LCMS:  $MH^+$  421.8.

Proceeding as in Example 2, but substituting 3-[(E)-3-[1-(2,4-difluoro-benzenesulfonyl)-1*H*-pyrrol-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-[1-(2,4-difluoro-benzenesulfonyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 28); LCMS:  $MH^+$  480.8.

Proceeding as in Example 2, but substituting 3-((E)-3-[2,2']bithienyl-5-yl-acryloyl)-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-(7-[2,2']bithienyl-5-yl-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one (Compound 29); LCMS: MH<sup>+</sup> 404.0.

Proceeding as in Example 2, but substituting 3-[(E)-3-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-{7-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one (Compound 30); LCMS: MH<sup>+</sup> 449.0.

Proceeding as in Example 2, but substituting 3-[(E)-3-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-{7-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one (Compound 31); LCMS: MH<sup>+</sup> 415.2.

Proceeding as in Example 2, but substituting 3-[(E)-3-(5-chloro-1*H*-indol-3-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(5-chloro-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 32); LCMS: MH<sup>+</sup> 389.0.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 33); LCMS: MH<sup>+</sup> 484.4.

Proceeding as in Example 2, but substituting 3-[(E)-3-(4,5-dibromo-thien-2-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-

methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(4,5-dibromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 34); LCMS:  $MH^+$  479.6.

Proceeding as in Example 2, but substituting 3-[(E)-3-(2-chloro-5-trifluoromethyl-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(2-chloro-5-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 35) 418.1.

Proceeding as in Example 2, but substituting 3-[(E)-3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 3-[3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one, provided 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one (Compound 36); MS  $M^+$  378.0; NMR ( $CDCl_3$ -TMS): d 13.00 (bs, 1H), 7.2 (d, 1H,  $J=9$  Hz), 6.45 (m, 2H), 4.36 (d, 1H,  $J=9$  Hz), 4.35 (m, 1H), 4.96-3.91 (m, 2H), 3.84 (s, 3H), 3.8 (m, 1H), 3.78 (s, 3H), 3.52 (dd, 1H,  $J=10$  Hz), 2.98 (m, 1H), 2.77 (m, 1H), 2.55-2.36 (m, 2H), 1.35 (d, 3H,  $J=6.5$  Hz).

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(5-methylsulfanyl-thien-2-yl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(5-methylsulfanyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 37); LCMS:  $MH^+$  367.4.

Proceeding as in Example 2, but substituting 3-[(E)-3-(5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-yl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 38); LCMS:  $MH^+$  430.0.

Proceeding as in Example 2, but substituting 2-[3-(2,4-dimethoxy-phenyl)-acryloyl]-3-hydroxy-cyclohex-2-enone for 3-[3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one, provided 2-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone (Compound 39); MS M<sup>+</sup> 361.8.

Proceeding as in Example 2, but substituting benzene-1,2-diamine for 2-aminoethanethiol, provided 3-[4-(2,4-dimethoxy-phenyl)-4,5-dihydro-3H-benzo[b][1,4]diazepin-2-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 40); LCMS: MH<sup>+</sup> 407.2.

Proceeding as in Example 2, but substituting 3-[(E)-3-(4-dimethylamino-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 42); MS M<sup>+</sup> 359.0.

Proceeding as in Example 2, but substituting 3-[(E)-3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-1H-quinolin-2-one for 3-[3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one, provided 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-1H-quinolin-2-one (Compound 43); MS M<sup>+</sup> 411.0.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(4-trifluoromethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 44); MS M<sup>+</sup> 399.8.

Proceeding as in Example 2, but substituting 3-[(E)-3-(bis-trifluoromethyl-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(bis-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 45); MS M<sup>+</sup> 452.0.

Proceeding as in Example 2, but substituting 3-[(E)-3-(4-dimethylamino-2-methoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(4-dimethylamino-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 46); MS M<sup>+</sup> 389.0.

Proceeding as in Example 2, but substituting 4-hydroxy-3-[(E)-3-(4-methanesulfonyl-phenyl)-acryloyl]-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-3-[7-(4-methanesulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one (Compound 47); MS M<sup>+</sup> 393.8.

Proceeding as in Example 2, but substituting 3-[(E)-3-(2,4-diethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one and 2-amino-benzenethiol for 2-aminoethanethiol, provided 3-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 48); MS M<sup>+</sup> 452.0.

Proceeding as in Example 2, but substituting 3-[(E)-3-(2,4-diethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one (Compound 49); MS M<sup>+</sup> 406.2.

Proceeding as in Example 2, but substituting 3-[(E)-3-(4-dimethylamino-phenyl)-acryloyl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one (Compound 50); MS M<sup>+</sup> 361.0.

Proceeding as in Example 2, but substituting 2-[(E)-3-(2,4-diethoxy-phenyl)-acryloyl]-3-hydroxy-cyclohex-2-enone for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 2-

[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone (Compound 51); MS  $M^+$  390.0.

Proceeding as in Example 2, but substituting 3-hydroxy-2-[(E)-3-(2,3,4-trimethoxy-phenyl)-acryloyl]-cyclohex-2-enone for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 3-hydroxy-2-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-cyclohex-2-enone (Compound 52); MS  $M^+$  392.2.

Proceeding as in Example 2, but substituting 2-[(E)-3-(2,4-diethoxy-phenyl)-acryloyl]-3-hydroxy-cyclohex-2-enone for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one and 2-amino-benzenethiol for 2-aminoethanethiol, provided 2-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-3-hydroxy-cyclohex-2-enone (Compound 53); MS  $M^+$  438.4.

Proceeding as in Example 2, but substituting 3-hydroxy-2-[(E)-3-(2,3,4-trimethoxy-phenyl)-acryloyl]-cyclohex-2-enone for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one and 2-amino-benzenethiol for 2-aminoethanethiol, provided 3-hydroxy-2-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone (Compound 54); MS  $M^+$  439.8.

Proceeding as in Example 2, but substituting 3-[3-(2,4-dimethoxy-phenyl)-acryloyl]-4-hydroxy-6-methyl-pyran-2-one for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one and 2-amino-3-mercapto-3-methyl-butyric acid for 2-aminoethanethiol, provided 7-(2,4-dimethoxy-phenyl)-5-(4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl)-2,2-dimethyl-2,3,6,7-tetrahydro-[1,4]thiazepine-3-carboxylic acid (Compound 55); MS  $M^+$  448.0.

Proceeding as in Example 2, but 4-hydroxy-6-methyl-3-[(E)-3-(2,3,4-trimethoxy-phenyl)-acryloyl]-5,6-dihydro-pyran-2-one substituted for 4-hydroxy-3-[(E)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one and 2-amino-benzenethiol for 2-aminoethanethiol, provided 4-hydroxy-6-methyl-




-84-

3-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-5,6-dihydro-pyran-2-one (Compound 56); MS  $M^+$  456.2.

Proceeding as in Example 2, but substituting 4-hydroxy-6-methyl-3-[(*E*)-3-(2,3,4-trimethoxy-phenyl)-acryloyl]-5,6-dihydro-pyran-2-one for 4-hydroxy-3-[(*E*)-3-(2-methoxy-4-methylsulfanyl-phenyl)-acryloyl]-6-methyl-pyran-2-one, provided 4-hydroxy-6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-5,6-dihydro-pyran-2-one (Compound 57); MS  $M^+$  408.4.

## EXAMPLE 3


3-[7-(2,4-Dimethoxy-phenyl)-1-oxo-2,3,6,7-tetrahydro-1*H*-1 $\lambda^4$ -[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one  
(Compound 58)



3-[7-(2,4-Dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (0.12 g, 0.32 mmol), prepared as in Reference 5, was dissolved in acetic acid (2 mL) and then hydrogen peroxide (0.3 ml, 35 wt % in water) was added at room temperature. The reaction was monitored by analytical HPLC. After 2 hours the solvent was removed *in vacuo*. Product was purified by preparative HPLC (RPC<sub>18</sub> column, 2-60 % acetonitrile/water containing 0.1% HCl) to provide a diastomeric mixture of 3-[7-(2,4-dimethoxy-phenyl)-1-oxo-2,3,6,7-tetrahydro-1*H*-1 $\lambda^4$ -[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (4.4 mg) as a yellow powder. LCMS:  $MH^+$  392.0.

## EXAMPLE 4

10-(2,4-Dimethoxy-phenyl)-3-methyl-7,8-dihydro-10H-2,5-dioxo-9-thia-6a-  
aza-cyclohepta[a]naphthalene-1,6-dione  
(Compound 59)




3-[7-(2,4-Dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-  
hydroxy-6-methyl-pyran-2-one (63 mg, 0.168 mmol), prepared as in Reference 5,  
was dissolved in ethylene dichloride (2 mL) and then 4-(dimethylamino)pyridine  
(0.020 g), di-*iso*-propylethylamine (0.5 mL) and phosgene (0.4 mL, 2 M in  
toluene) were added at room temperature. The reaction was monitored by  
analytical HPLC. After stirring at room temperature for 30 minutes, the solvent  
was removed *in vacuo*. Product was purified by HPLC (RPC<sub>18</sub> column, 2-70 %  
acetonitrile/water containing 0.1% HCl) to provide 10-(2,4-dimethoxy-phenyl)-3-  
methyl-7,8-dihydro-10H-2,5-dioxo-9-thia-6a-aza-cyclohepta[a]naphthalene-1,6-  
dione (20 mg) as an off-white powder. LCMS: M<sup>+</sup> 401.0.

## EXAMPLE 5

3-[7-(2,4-Dimethoxy-phenyl)-1,1-dioxo-2,3,6,7-tetrahydro-1H-  
1λ<sup>6</sup>[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one  
(Compound 60)


-86-



3-[7-(2,4-Dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (190 mg, 0.5 mmol), prepared as in Reference 5, was dissolved in acetic acid (2 ml) and hydrogen peroxide (0.3 ml, 35 wt % in water) was added to the solution at room temperature. The clear solution was stirred at 70°C. The reaction was monitored by analytical HPLC. After 2 hours the solvent was removed *in vacuo*. Product was purified by preparative HPLC (RPC<sub>18</sub> column, 2-60 % acetonitrile/water containing 0.1 % HCl) to provide 3-[7-(2,4-dimethoxy-phenyl)-1,1-dioxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>6</sup>[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (10 mg) as a yellow powder. LCMS: MH<sup>+</sup> 408.0.

## EXAMPLE 6

3-[7-(2,4-Dimethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one  
(Compound 61)



3-[7-(2,4-Dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (150 mg, 0.39 mmol), prepared as in Reference 5, was dissolved in toluene (2 ml) and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (0.3 mL, 35 wt % in water) was added to the solution. The mixture was stirred at 80°C. The reaction was monitored by analytical HPLC and after reaction was complete (20 minutes) the solvent was then removed *in vacuo*. Product was purified by preparative HPLC (RPC<sub>18</sub> column, 2-80 % acetonitrile/water containing 0.1 % HCl) to provide 3-[7-(2,4-dimethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (35 mg) as an orange powder. LCMS: MH<sup>+</sup> 373.0.

Proceeding as in Example 6, but substituting 3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one for 3-[7-(2,4-dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one, provided 3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 62); MS M<sup>+</sup> 402.0.

Proceeding as in Example 6, but substituting 3-(7-[2,2']bithienyl-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one for 3-[7-(2,4-dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one, provided 3-(7-[2,2']bithienyl-5-yl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 63); MS M<sup>+</sup> 402.2.


Proceeding as in Example 6, but substituting 2-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone for 3-[7-(2,4-dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one, provided 2-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone (Compound 64); MS M<sup>+</sup> 388.4.

Proceeding as in Example 6, but substituting 3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one for 3-[7-(2,4-dimethoxyphenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-

hydroxy-6-methyl-pyran-2-one, provided 3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one (Compound 65); MS  $M^+$  404.4.

## EXAMPLE 7

5 2-((1-[7-(2,4-Dimethoxy-phenyl)-5-(4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl)-2,2-dimethyl-2,3,6,7-tetrahydro-[1,4]thiazepin-3-yl]-methanoyl)-amino)-propionic acid *tert*-butyl ester  
(Compound 66)



10 7-(2,4-Dimethoxy-phenyl)-5-(4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl)-2,2-dimethyl-2,3,6,7-tetrahydro-[1,4]thiazepine-3-carboxylic acid (0.03 g), prepared as in Example 2, was dissolved in DMF (2 mL) and 2-amino-propionic acid *tert*-butyl ester (0.012 g), *N*-methylmorpholine (0.2 ml) and PyBOP (0.034 g) were added to the solution at room temperature. The mixture was stirred while  
15 the progress of the reaction was followed by analytical HPLC and when complete (16 hours) the solvent was removed *in vacuo*. Purification of product by preparative HPLC (RPC<sub>18</sub> column, 2-70 % acetonitrile/water containing 0.1% HCl) provided 2-((1-[7-(2,4-dimethoxy-phenyl)-5-(4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl)-2,2-dimethyl-2,3,6,7-tetrahydro-[1,4]thiazepin-3-yl]-methanoyl)-

amino)-propionic acid *tert*-butyl ester (18 mg) as an off-white powder. LCMS:  $M^+$  575.0.

Proceeding by methods analogous to those described in the application the following compounds can be made:

4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 70),

3-[7-(4-ethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 71),

**4-hydroxy-3-[7-(3-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one (Compound 72),**

3-[7-(2-bromo-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 74),

**3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 79),**

**3-[7-(2,3-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one (Compound 82).**

**4-hydroxy-6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one (Compound 83),**

4-hydroxy-6-methyl-3-(2-*p*-tolyl-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl)-pyran-2-one (Compound 85) and

**4-hydroxy-6-methyl-3-[2-(4-methylsulfanyl-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-pyran-2-one (Compound 86).**

### EXAMPLE 8

## Identification of Caspase Cascade Activators and Inducers of Apoptosis in Solid Tumor Cells

Human breast cancer cell lines T-47D and ZR-75-1 were grown according to media component mixtures designated by American Type Culture Collection + 10% fetal calf sera (FCS) (Life Technologies, Inc.) in a 5% CO<sub>2</sub>-95% humidity

-90-

incubator as 37 °C. The T-47 and ZR-75-1 cells were maintained at a cell density between 30 and 80% confluency at a cell density of 0.1 to 0.6 x 10<sup>6</sup> cells/mL. Cells were harvested at 600xg and resuspended at 0.65 x 10<sup>6</sup> cells/mL into appropriate media + 10% FCS. An aliquot of 45 µL of cells was added to a well of a 96-well microtiter plate containing 5 µL of a 10% DMSO in RPMI-1640 media solution containing 1.6 to 100 µM of test compound (0.16 to 10 µM final). An aliquot of 45 µL of cells was added to a well of a 96-well microtiter plate containing 5 µL of a 10% DMSO in RPMI-1640 media solution without test compound as the control sample. The samples were mixed by agitation and then incubated at 37 °C for 24 hours in a 5% CO<sub>2</sub>-95% humidity incubator. After incubation, the samples were removed from the incubator and 50 µL of a solution containing 20 µL of *N*-(Ac-DEVD)-*N'*-ethoxycarbonyl-R110 (SEQ ID NO:1) fluorogenic substrate (Cytovia, Inc.; WO99/18856), 20% sucrose (Sigma), 20 mM dithiothreitol (DTT) (Sigma), 200 mM NaCl (Sigma), 40 mM Na piperazine-*N,N'*-bis[2-ethanesulfonic acid] (PIPES) buffer pH 7.2 (Sigma), and 500 µg/mL lysolecithin (Calbiochem) was added. The samples were mixed by agitation and incubated at room temperature. Using a fluorescent plate reader (Model 1420 Wallac Instruments), an initial reading (T = 0) was made approximately 1-2 minutes after addition of the substrate solution, employing excitation at 485 nm and emission at 530 nm, to determine the background fluorescence of the control sample. After the 3 hour incubation, the samples were read for fluorescence as above (T = 3 hours).

#### Calculation:

The Relative Fluorescence Unit (RFU) values were used to calculate the sample readings as follows:

$$\text{RFU}_{(T=3h)} - \text{Control RFU}_{(T=0)} = \text{Net RFU}_{(T=3h)}$$

The level of caspase cascade activation was determined by the ratio of the net RFU value for the test compound to that of the control samples. The EC<sub>50</sub> (nM) was determined by a sigmoidal dose-response calculation (Prism 2.0,

GraphPad Software, Inc.). The compounds of the invention were determined to have caspase cascade activating effects by proceeding as in Example 8.

**Table I. Caspase Potency**

Compound #	EC <sub>50</sub> (nM)	
	T-47D	ZR-75-1
70	345	163
72	3050	1950
74	3270	2080
79	557	349
85	6930	4207

### EXAMPLE 9

#### Identification of Antineoplastic Activity in Cell Proliferation

T-47D and ZR-75-1 cells are grown and harvested by proceeding as in Example 8. An aliquot of 90  $\mu$ L of cells ( $2.2 \times 10^4$  cells/mL) is added to a well of a 96-well microtiter plate containing 10  $\mu$ L of a 10% DMSO in RPMI-1640 media solution containing 1 nM to 100  $\mu$ M of test compound. An aliquot of 90  $\mu$ L of cells is added to a well of a 96-well microtiter plate containing 10  $\mu$ L of a 10% DMSO in RPMI-1640 media solution without test compound as the control sample for maximal cell proliferation ( $A_{\max}$ ). The samples are mixed by agitation and then incubated at 37°C for 48 hours in a 5% CO<sub>2</sub>-95% humidity incubator. After incubation, the samples are removed from the incubator and 20  $\mu$ L of CellTiter 96 Aqueous One Solution Cell Proliferation™ reagent (Promega) is added. The samples are mixed by agitation and incubated at 37°C for 2-4 hours in a 5% CO<sub>2</sub>-95% humidity incubator. Using an absorbance plate reader (Model 1420 Wallac Instruments), an initial reading (T=0) is made approximately 1-2 minutes after addition of the solution, employing absorbance at 490 nm, to



determine any background absorbance of the test compound. After the 2-4 hours incubation, the samples are read for absorbance as above ( $A_{\text{test}}$ ).

Baseline for the dose producing 50% inhibition of cell proliferation ( $GI_{50}$ ) of initial cell numbers is determined by adding an aliquot of 90  $\mu\text{L}$  of cells or 90  $\mu\text{L}$  of media, respectively, to wells of a 96-well microtiter plate containing 10  $\mu\text{L}$  of a 10% DMSO in RPMI-1640 media solution. The samples are mixed by agitation and then incubated at 37°C for 0.5 hours in a 5%  $\text{CO}_2$ -95% humidity incubator. After incubation, the samples are removed from the incubator and 20  $\mu\text{L}$  of CellTiter 96 Aqueous One Solution Cell Proliferation™ reagent (Promega) is added. The samples are mixed by agitation and incubated at 37°C for 2-4 hours in a 5%  $\text{CO}_2$ -95% humidity incubator. Absorbance is read as above, ( $A_{T=0}$ ) defining absorbance for initial cell number used as baseline  $GI_{50}$  determinations.

Calculation:

$$GI_{50} \text{ (nM)} = 100 \times [A_{\text{test}} - A_{T=0}] / (A_{\text{max}} - A_{T=0}).$$

#### EXAMPLE 10

##### Nuclear Fragmentation in T47D Cells

T47D cells are grown and harvested by proceeding as in Example 8 and treated with test compound followed by staining of the cell nuclei with Syto16, a fluorescent DNA dye which stains nuclei. Shrunken and fragmented nuclei are hallmarks of caspase-mediated apoptosis. T47D cells treated with test compound for 48 hours exhibit shrunken and fragmented nuclei.

#### EXAMPLE 11

##### Mitotic Arrest in Jurkat Cells

Jurkat cells are incubated with a range of concentrations of test compounds (0.02  $\mu\text{M}$  to 5  $\mu\text{M}$ ) for 6 hours under normal growth conditions. Control cultures are treated with DMSO vehicle. The cells are then treated for

20 minutes with 800 nM Syto 16. Cytospin preparation are then prepared and the samples were viewed by fluorescent microscopy using a fluorescein filter set. For each concentration of test compound, the number of mitotic figures are counted and expressed as a percentage of the total number of cells. Three fields from each condition are evaluated and the mean and SEM were calculated and plotted as a function of drug concentration.

## EXAMPLE 12


### Cell Cycle Arrest in Solid Tumor Cell Lines

T47D cells are grown and harvested by proceeding as in Example 8. Cells at  $1 \times 10^6$  are treated with test compound for 48 hours at 37°C. As a control, cells are also incubated with DMSO. Cells were harvested at 1200 rpm and washed twice with 5 mM EDTA/PBS. Cells are then resuspended in 300  $\mu$ L of EDTA/PBS and 700 mL of 100% ethanol, vortexed and incubated at room temperature for 1 hour. Samples are spun down at 12000 rpm for 5 minutes and the supernatant is removed. A solution containing 100  $\mu$ g/mL of propidium iodide and 1 mg/mL of RNase A (fresh) is added to the samples and the samples are incubated for 1 hour at room temperature. Samples are then transferred to 12x75 mm polystyrene tubes and analyzed on a flow cytometer. All flow cytometry analyses are performed on FACScalibur (Becton Dickison) using Cell Quest analysis software.

Having now fully described this invention, it will be understood by those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any embodiment thereof. All patents, patent applications and publications cited herein are fully incorporated by reference herein in their entirety.

**What Is Claimed Is:**

1. A compound of Formula I:



in which:

the dashed lines indicate optional unsaturation without violating valency rules;

$R^1$  is hydrogen,  $(C_{1-6})$ alkyl or  $-C(O)R^6$ , wherein  $R^6$  is as defined below, or  $R^1$  is absent when a double bond exists between the nitrogen atom to which  $R^1$  is attached and an adjacent ring atom or  $R^1$  is as defined below;

$X^1$  is  $-NR^2$ -,  $-S$ -,  $-S(O)$ -,  $-S(O)_2$ - or  $-O$ -, wherein  $R^2$  is hydrogen or  $(C_{1-6})$ alkyl or is absent when a double bond exists between the nitrogen atom to which  $R^2$  is attached and an adjacent ring atom;

$A^1$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, or  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a fused polycyclic ring system selected from heteroaryl and unsaturated, partially unsaturated or saturated heterocycloalkyl in any case containing a total of 10 to 15 ring atoms, wherein  $A^1$  may be substituted with a group selected from  $-X^2R^3$ ,  $-X^2OR^3$ ,  $-X^2C(O)R^3$ ,  $-X^2OC(O)R^3$ ,  $-X^2C(O)OR^3$ ,  $-X^2SR^3$ ,  $-X^2S(O)R^3$ ,  $-X^2S(O)_2R^3$ ,  $-X^2NR^3R^4$ ,  $-X^2NR^4C(O)R^3$ ,  $-X^2NR^4C(O)OR^3$ ,  $-X^2C(O)NR^3R^4$ ,  $-X^2NR^4C(O)NR^3R^4$ ,  $-X^2NR^4C(NR^4)NR^3R^4$ ,  $-X^2NR^4S(O)_2R^3$  and  $-X^2S(O)_2NR^3R^4$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^3$  is  $-X^2R^5$  wherein  $X^2$  is as defined above and  $R^5$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10


ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^1$  and  $R^5$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within  $A^1$  and  $R^5$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^1$  and  $R^5$  is a fused polycyclic ring system;

$A^2$  is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein  $A^2$  may be substituted with a group selected from  $-X^2R^8$ ,  $-X^2OR^8$ ,  $-X^2C(O)R^8$ ,  $-X^2OC(O)R^8$ ,  $-X^2C(O)OR^8$ ,  $-X^2SR^8$ ,  $-X^2S(O)R^8$ ,  $-X^2S(O)_2R^8$ ,  $-X^2NR^4R^8$ ,  $-X^2NR^4C(O)R^8$ ,  $-X^2NR^4C(O)OR^8$ ,  $-X^2C(O)NR^4R^8$ ,  $-X^2NR^4C(O)NR^4R^8$ ,  $-X^2NR^4C(NR^4)NR^4R^8$ ,  $-X^2NR^4S(O)_2R^8$  and  $-X^2S(O)_2NR^4R^8$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$  is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^2$  and  $R^8$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,

$-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  
 $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  
 $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said  
 heterocycloalkylene, carbocycloalkyl and heterocycloalkyl rings within  $A^2$  and  $R^8$   
 may be substituted further with 1 to 2 groups independently selected from  
 (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^2$  and  
 $R^8$  is a fused polycyclic ring system; and

$A^3$  is a monocyclic or fused polycyclic ring system selected from aryl  
 containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14  
 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or  
 heterocycloalkyl each containing a total of 3 to 14 ring atoms, wherein  $A^3$  may  
 be substituted with a group selected from  $-X^2R^9$ ,  $-X^2OR^9$ ,  $-X^2C(O)R^9$ ,  
 $-X^2OC(O)R^9$ ,  $-X^2C(O)OR^9$ ,  $-X^2SR^9$ ,  $-X^2S(O)R^9$ ,  $-X^2S(O)_2R^9$ ,  $-X^2NR^4R^9$ ,  
 $-X^2NR^4C(O)R^9$ ,  $-X^2NR^4C(O)OR^9$ ,  $-X^2C(O)NR^4R^9$ ,  $-X^2NR^4C(O)NR^4R^9$ ,  
 $-X^2NR^4C(NR^4)NR^4R^9$ ,  $-X^2NR^4S(O)_2R^9$  and  $-X^2S(O)_2NR^4R^9$ , wherein  $X^2$  is a bond  
 or  $(C_{1-6})$ alkylene,  $R^9$  is  $-X^2R^{10}$  wherein  $X^2$  is as defined above and  $R^{10}$  is aryl  
 containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10  
 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or  
 heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each  
 occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl,  
 wherein each ring within  $A^3$  and  $R^{10}$  contains from 3 to 8 ring atoms and may be  
 substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano,  
 halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  
 $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  
 $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  
 $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  
 $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said  
 carbocycloalkyl and heterocycloalkyl rings within  $A^3$  and  $R^{10}$  may be substituted  
 further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo,  
 imino and thioxo, with the proviso that only one of  $A^3$  and  $R^{10}$  is a fused

polycyclic ring system; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof; with the proviso that when said compound is selected from the group consisting of Formulae II(a), II(b) and II(c):



then A<sup>3</sup> is other than:

- unsubstituted pyridyl;
- unsubstituted thienyl;
- unsubstituted indolyl;
- unsubstituted phenyl;
- benzo[1,3]dioxolyl;
- 2,3-dihydro-benzo[1,4]dioxinyl;
- phenyl which is mono-substituted by fluoro, bromo, iodo, nitro, methyl, isopropyl, ethoxy or methylsulfanyl; and
- phenyl which is substituted by at least one of chloro, hydroxy or methoxy.


2. The compound of claim 1, and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts of said compound, with the further proviso that A<sup>3</sup> is other than:

- unsubstituted pyridyl;
- unsubstituted thienyl;

unsubstituted indolyl;  
unsubstituted phenyl;  
benzo[1,3]dioxolyl;  
2,3-dihydro-benzo[1,4]dioxinyl; and  
5 phenyl which is substituted by at least one of halogen, nitro, hydroxy, (C<sub>1-3</sub>)alkyl, methoxy, ethoxy and methylsulfanyl.

3. The compound of claim 1, and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts of said compound, with  
10 the further proviso that A<sup>1</sup> is not 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl.

4. The compound of Claim 1 in which said compound is of Formula I(A):




I(A)

15 in which R<sup>1</sup>, A<sup>1</sup>, A<sup>2</sup> and A<sup>3</sup> are as defined in Claim 1; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

5. The compound of Claim 4 in which said compound is of Formula I(B):


-99-



I(B)

in which  $R^1$ ,  $A^1$ ,  $A^2$  and  $A^3$  are defined as in Claim 1; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

- 5                    6.    The compound of Claim 5 in which said  $A^2$  is 2,3,6,7-tetrahydro-[1,4]thiazepin-5,7-ylene, that is the compound of Formula I(C):



I(C)

- 10                    in which  $A^1$  and  $A^3$  are defined as in Claim 1, and said 2,3,6,7-tetrahydro-[1,4]thiazepin-5,7-ylene may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.
- 15



7. The compound of Claim 6 in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

8. The compound of Claim 7 in which said compound is selected from the group consisting of:

4-hydroxy-6-methyl-3-[7-(3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(5-ethyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(1-benzyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3[7-(2-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3[7-(3-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3[7-(4-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-[3-(3-trifluoromethyl-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-[3-(3,4-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-{7-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

3-{7-[5-(2-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[5-(3-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[5-(4-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

5 4-hydroxy-6-methyl-3-{7-[5-(2-chloro-5-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

3-{7-(4-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

10 3-{7-(5-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-(1-benzenesulfonyl-1*H*-pyrrol-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-{7-(3-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

15 4-hydroxy-6-methyl-3-{7-(5-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

4-hydroxy-6-methyl-3-{7-(1-methyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

20 3-{7-(3-chloro-2-methyl-5-trifluoromethyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[1-(2,4-difluoro-benzenesulfonyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one;

25 3-{7-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

30 3-{7-(5-chloro-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(4,5-dibromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 4-hydroxy-6-methyl-3-[7-(5-methylsulfanyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(5-chloro-1-methyl-3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(4-trifluoromethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(bis-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

15 4-hydroxy-3-[7-(4-methanesulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; and

3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-methoxy-6-methyl-pyran-2-one;

20 and the *N*-oxide derivatives, prodrug derivatives, protected derivatives; and the pharmaceutically acceptable salts thereof.

9. The compound of Claim 6 in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

10. The compound of Claim 9 in which said compound is selected from the group consisting of:

3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

5           3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one; and

3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

10           and the *N*-oxide derivatives, prodrug derivatives, protected derivatives; and the pharmaceutically acceptable salts thereof.

11.   The compound of Claim 6 in which A<sup>1</sup> is 2-hydroxy-6-oxo-cyclohex-1-enyl or 2-methoxy-6-oxo-cyclohex-1-enyl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

15           12.   The compound of Claim 11 in which said compound is selected from the group consisting of:


2-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

20           2-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone; and

3-hydroxy-2-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-cyclohex-2-enone;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives; and the pharmaceutically acceptable salts thereof.

13. The compound of claim 6 in which A<sup>1</sup> is a group of Formula (c):



(c)


in which R<sup>7</sup> is hydrogen or methyl, R<sup>11</sup> is hydrogen or (C<sub>1-6</sub>)alkyl and the free valence is attached to A<sup>2</sup>; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

14. The compound of Claim 13 which is:

3-[7-2,4-dimethoxy-phenyl)-2,3,6,,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-1*H*-quinolin-2-one;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives; and the pharmaceutically acceptable salts thereof.

15. The compound of Claim 5 in which said A<sup>2</sup> is 2,3-dihydro-benzo[*b*][1,4]thiazepin-5,7-ylene that is the compound of Formula I(D):



in which A<sup>1</sup> and A<sup>3</sup> are defined as in Claim 1, and said 2,3-dihydro-benzo[*b*][1,4]thiazepin-5,7-ylene may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>,  
5 -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>,  
-X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>,  
wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl; and the *N*-oxide derivatives, prodrug derivatives,  
10 protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

16. The compound of Claim 15 in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual  
15 stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

17. The compound of Claim 16 which is:  
3-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-4-hydroxy-6-methyl-pyran-2-one;  
20 and the *N*-oxide derivatives, prodrug derivatives, protected derivatives; and the pharmaceutically acceptable salts thereof.

18. The compound of Claim 15 in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl; and the *N*-oxide derivatives, prodrug derivatives, protected  
25 derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

19. The compound of Claim 18 which is:

4-hydroxy-6-methyl-3-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-5,6-dihydro-pyran-2-one;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives;  
and the pharmaceutically acceptable salts thereof.

20. The compound of Claim 15 in which A<sup>1</sup> is 2-hydroxy-6-oxo-cyclohex-1-enyl or 2-methoxy-6-oxo-cyclohex-1-enyl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.


21. The compound of Claim 20 in which said compound is selected from the group consisting of:

3-hydroxy-2-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone; and

3-hydroxy-2-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives;  
and the pharmaceutically acceptable salts thereof.

22. The compound of Claim 4 in which said A<sup>2</sup> is a group of Formula (k):



(k)

in which R<sup>1</sup> is defined as in Claim 1 and said group of Formula (k) may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>,

-X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>,  
-X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a  
bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted  
(C<sub>1-6</sub>)alkyl; and the *N*-oxide derivatives, prodrug derivatives, protected  
derivatives, individual stereoisomers and mixtures of stereoisomers; and the  
pharmaceutically acceptable salts thereof.

23. The compound of Claim 22 in which R<sup>1</sup> is hydrogen; and the  
*N*-oxide derivatives, prodrug derivatives, protected derivatives, individual  
stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable  
salts thereof.

24. The compound of Claim 22 in which A<sup>1</sup> is 4-hydroxy-6-methyl-  
2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and the  
*N*-oxide derivatives, prodrug derivatives, protected derivatives, individual  
stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable  
salts thereof.

25. The compound of Claim 24 in which said compound is selected  
from the group consisting of:

3-[4-acetyl-7-(2,4-dimethoxy-phenyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-  
methyl-pyran-2-one; and

3-[7-(2,4-dimethoxy-phenyl)-4-(2,2,2-trifluoro-ethanoyl)-[1,4]thiazepan-  
5-yl]-4-hydroxy-6-methyl-pyran-2-one;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives;  
and the pharmaceutically acceptable salts thereof.

26. The compound of Claim 22 in which A<sup>1</sup> is optionally substituted  
phenyl.




-108-

27. The compound of Claim 26 which is:

1-[7-(2,4-dimethoxy-phenyl)-5-(3-fluoro-4-methoxyphenyl)-  
[1,4]thiazepan-4-yl]-ethanone;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives;  
and the pharmaceutically acceptable salts thereof.

28. The compound of Claim 4 in which said A<sup>2</sup> is 2,3-dihydro-  
[1,4]thiazepin-5,7-ylene that is the compound of Formula I(F):



I(F)

in which A<sup>1</sup> and A<sup>3</sup> are defined as in Claim 1, and said 2,3-dihydro-  
[1,4]thiazepin-5,7-ylene may be substituted with 1 to 3 groups independently  
selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>,  
-X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>,  
-X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>,  
-X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>,  
-X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup>  
at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)  
alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl; and the *N*-oxide  
derivatives, prodrug derivatives, protected derivatives, individual stereoisomers  
and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

29. The compound of Claim 28 in which A<sup>1</sup> is 4-hydroxy-6-methyl-  
2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and the  
*N*-oxide derivatives, prodrug derivatives, protected derivatives, individual  
stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable  
salts thereof.

30. The compound of Claim 29 in which said compound is selected from the group consisting of:

3-[7-(2,4-dimethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one; and

3-(7-[2,2']bithienyl-5-yl)-2,3-dihydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives;  
10 and the pharmaceutically acceptable salts thereof.

31. The compound of Claim 28 in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-5,6-dihydro-2*H*-pyran-3-yl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the  
15 pharmaceutically acceptable salts thereof.

32. The compound of Claim 31 which is:

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives;  
20 and the pharmaceutically acceptable salts thereof.

33. The compound of Claim 28 in which A<sup>1</sup> is 2-hydroxy-6-oxo-cyclohex-1-enyl or 2-methoxy-6-oxo-cyclohex-1-enyl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.


-110-

34. The compound of Claim 33 which is:

2-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.


35. The compound of Claim 1 in which said compound is of Formula I(G):



I(G)

in which *n*, *R*<sup>1</sup>, *A*<sup>1</sup>, *A*<sup>2</sup> and *A*<sup>3</sup> are defined as in Claim 1; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

36. The compound of Claim 35 in which *A*<sup>2</sup> is a group of Formula (I):



(I)

in which *n*, and *R*<sup>1</sup> are defined as in Claim 1 and said group of Formula (I) may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> is a

bond or (C<sub>1-6</sub>)alkylene, R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

37. The compound of Claim 36 in which *n* is 1 and A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.


38. The compound of Claim 37 which is:  
3-[7-(2,4-dimethoxy-phenyl)-1-oxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>4</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methoxy-pyran-2-one;  
and the *N*-oxide derivatives, prodrug derivatives, protected derivatives; and the pharmaceutically acceptable salts thereof.

39. The compound of claim 36 in which *n* is 2 and A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

40. The compound of claim 39 which is:  
3-[7-(2,4-dimethoxy-phenyl)-1,1-dioxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>6</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;  
and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

-112-


41. The compound of Claim 1 in which said compound is of Formula I(I):



I(I)

in which  $R^1$ ,  $R^2$ ,  $A^1$ ,  $A^2$  and  $A^3$  are as defined in Claim 1; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

42. The compound of Claim 41 in which said  $A^2$  is a group of Formula (j):




(j)

in which said group of Formula (j) may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

43. The compound of Claim 42 in which A<sup>1</sup> is 4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl or 4-methoxy-6-methyl-2-oxo-2*H*-pyran-3-yl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

44. The compound of Claim 43 which is:  
3 - [ 4 - ( 2 , 4 - d i m e t h o x y - p h e n y l ) - 4 , 5 - d i h y d r o - 3 H - benzo[*b*][1,4]diazepin-2-yl]-4-hydroxy-6-methylpyran-2-one;  
and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

45. The compound of Claim 1 in which said compound is of Formula I(K):



I(K)

in which A<sup>3</sup> is defined as in Claim 1; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

46. The compound of claim 45 which is:  
10-(2,4-dimethoxy-phenyl)-3-methyl-7,8-dihydro-10*H*-2,5-dioxa-9-thia-6*a*-aza-cyclohepta[*a*]naphthalene-1,6-dione;

and the *N*-oxide derivatives, prodrug derivatives, protected derivatives; and the pharmaceutically acceptable salts thereof.

47. A compound selected from the group consisting of:

4-hydroxy-3-[7-(2-methoxy-4-methylsulfonyl-phenyl)-2,3,6,7-tetrahydro-  
[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[7-(2-chloro-5-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-  
[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-2-methoxy-phenyl)-2,3,6,7-tetrahydro-  
[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-3-[7-(4-chloro-2-methoxy-phenyl)-2,3,6,7-tetrahydro-  
[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; and

4-hydroxy-3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-  
yl]-6-methyl-pyran-2-one; or

a *N*-oxide derivative, prodrug derivative, protected derivative, individual  
stereoisomer and mixtures of stereoisomers; or the pharmaceutically acceptable  
salt thereof.

48. A compound selected from the group consisting of:


7-(2,4-dimethoxy-phenyl)-5-(4-hydroxy-6-methyl-2-oxo-2*H*-pyran-3-yl)-  
2,2-dimethyl-2,3,6,7-tetrahydro-[1,4]thiazepine-3-carboxylic acid; and

2-({1-[7-(2,4-dimethoxy-phenyl)-5-(4-hydroxy-6-methyl-2-oxo-  
2*H*-pyran-3-yl)-2,2-dimethyl-2,3,6,7-tetrahydro-[1,4]thiazepin-3-yl]-methanoyl}-  
amino)-propionic acid *tert*-butyl ester;


and the *N*-oxide derivatives, prodrug derivatives, protected derivatives;  
and the pharmaceutically acceptable salts thereof..

49. The compound of Claim 1 in which A<sup>1</sup> is a group selected from  
Formulae (b), (c), (d), (e) and (f):


-115-




(b)




(c)



(d)



(e)




(f)

in which  $R^7$  is hydrogen or methyl,  $R^{11}$  is hydrogen or  $(C_{1-6})$ alkyl and the free valance is attached to  $A^2$ , or

$A^2$  and  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a group of Formula (g):

5



(g)

wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ ; and

$A^2$  is as defined above or is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein  $A^2$  may be substituted with a group selected from  $-X^2R^8$ ,  $-X^2OR^8$ ,  $-X^2C(O)R^8$ ,  $-X^2OC(O)R^8$ ,


10




$-X^2C(O)OR^8$ ,  $-X^2SR^8$ ,  $-X^2S(O)R^8$ ,  $-X^2S(O)_2R^8$ ,  $-X^2NR^4R^8$ ,  $-X^2NR^4C(O)R^8$ ,  
 $-X^2NR^4C(O)OR^8$ ,  $-X^2C(O)NR^4R^8$ ,  $-X^2NR^4C(O)NR^4R^8$ ,  $-X^2NR^4C(NR^4)NR^4R^8$ ,  
 $-X^2NR^4S(O)_2R^8$  and  $-X^2S(O)_2NR^4R^8$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$   
 5 is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to  
 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated,  
 partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each  
 containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently  
 is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  
 10  $A^2$  and  $R^8$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3  
 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro,  
 halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  
 $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  
 $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  
 $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  
 15  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and  
 wherein any said heteroalkylene, carbocycloalkyl and heterocycloalkyl rings  
 within  $A^2$  and  $R^8$  may be substituted further with 1 to 2 groups independently  
 selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only  
 one of  $A^2$  and  $R^8$  is a fused polycyclic ring system; and the *N*-oxide derivatives,  
 20 prodrug derivatives, protected derivatives, individual stereoisomers and mixtures  
 of stereoisomers; and the pharmaceutically acceptable salts thereof.

50. The compound of Claim 49 in which  $A^2$  is a group selected from  
 Formulae (h), (i), (j), (k), (l) and (m):


-117-




(h)




(i)




(j)



(k)




(l)



(m)

in which  $n$  is 1 or 2 and  $R^1$  is acetyl or trifluoroacetyl or  $A^2$  and  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a group of Formula (g):




(g)

- 5 wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ ; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

51. A compound of Formula II:

-118-



in which:

the dashed lines indicate optional unsaturation without violating valency rules;

$R^1$  is hydrogen,  $(C_{1-6})$ alkyl or  $-C(O)R^6$ , wherein  $R^6$  is as defined below, or  $R^1$  is absent when a double bond exists between the nitrogen atom to which  $R^1$  is attached and an adjacent ring atom or  $R^1$  is as defined below;

$R^7$  is hydrogen;

$X^1$  is  $-NR^2$ -,  $-S$ -,  $-S(O)$ -,  $-S(O)_2$ - or  $-O$ -, wherein  $R^2$  is hydrogen or  $(C_{1-6})$ alkyl or is absent when a double bond exists between the nitrogen atom to which  $R^2$  is attached and an adjacent ring atom;

$A^2$  is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein  $A^2$  may be substituted with a group selected from  $-R^8$ ,  $-X^2OR^8$ ,  $-X^2C(O)R^8$ ,  $-X^2OC(O)R^8$ ,  $-X^2C(O)OR^8$ ,  $-X^2SR^8$ ,  $-X^2S(O)R^8$ ,  $-X^2S(O)_2R^8$ ,  $-X^2NR^4R^8$ ,  $-X^2NR^4C(O)R^8$ ,  $-X^2NR^4C(O)OR^8$ ,  $-X^2C(O)NR^4R^8$ ,  $-X^2NR^4C(O)NR^4R^8$ ,  $-X^2NR^4C(NR^4)NR^4R^8$ ,  $-X^2NR^4S(O)_2R^8$  and  $-X^2S(O)_2NR^4R^8$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$  is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^2$  and  $R^8$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,


-X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>,  
 -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>,  
 -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and  
 -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or  
 5 halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said heteroalkylene, carbocycloalkyl  
 and heterocycloalkyl rings within A<sup>2</sup> and R<sup>8</sup> may be substituted further with 1 to  
 2 groups independently selected from (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo  
 with the proviso that only one of A<sup>2</sup> and R<sup>8</sup> is a fused polycyclic ring system; and

A<sup>3</sup> is a monocyclic or fused polycyclic ring system selected from aryl  
 10 containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14  
 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or  
 heterocycloalkyl each containing a total of 3 to 14 ring atoms, wherein A<sup>3</sup> may  
 be substituted with a group selected from -R<sup>9</sup>, -X<sup>2</sup>OR<sup>9</sup>, -X<sup>2</sup>C(O)R<sup>9</sup>, -X<sup>2</sup>OC(O)R<sup>9</sup>,  
 -X<sup>2</sup>C(O)OR<sup>9</sup>, -X<sup>2</sup>SR<sup>9</sup>, -X<sup>2</sup>S(O)R<sup>9</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>9</sup>,  
 15 -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>9</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>9</sup>,  
 -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>9</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>9</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>9</sup>  
 is -X<sup>2</sup>R<sup>10</sup> wherein X<sup>2</sup> is as defined above and R<sup>10</sup> is aryl containing a total of 6 to  
 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated,  
 partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each  
 20 containing a total of 3 to 10 ring atoms, and R<sup>4</sup> at each occurrence independently  
 is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, wherein each ring within  
 A<sup>3</sup> and R<sup>10</sup> contains from 3 to 8 ring atoms and may be substituted with 1 to 3  
 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro,  
 halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>,  
 25 -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>,  
 -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and  
 -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or  
 halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said carbocycloalkyl and  
 heterocycloalkyl rings within A<sup>3</sup> and R<sup>10</sup> may be substituted further with 1 to 2  
 30 groups independently selected from (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo with


the proviso that only one of  $A^3$  and  $R^{10}$  is a fused polycyclic ring system; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof;

provided, however, Formula II does not represent a compound wherein  $A^2$  is 2,3,6,7-tetrahydro-[1,4]thiazepinylene, 2,3-dihydrobenzo[*b*][1,4]thiazepinylene or 7-trifluoro-2,3-dihydrobenzo[*b*][1,4]thiazepinylene and  $A^3$  is benzo[1,3]dioxolyl, indolyl, phenyl, pyridyl or thienyl, wherein said phenyl may be substituted with 1 to 3 groups independently selected from halo, nitro, hydroxy,  $(C_{1-4})$ alkyl,  $(C_{1-4})$ alkylsulfanyl and  $(C_{1-4})$ alkyloxy or any *N*-oxide derivative; protected derivative, individual stereoisomer or mixture of stereoisomers, or pharmaceutically acceptable salt thereof.


52. The compound of Claim 51 in which  $A^2$  is a group selected from Formulae (h), (i), (j), (k), (l) and (m):




(h)




(i)




(j)



(k)




(l)



(m)

-121-

in which  $n$  is 1 or 2 and  $R^1$  is acetyl or trifluoroacetyl or  $A^2$  and  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a group of Formula (g):



(g)

wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ ; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.


53. The compound of Claim 52 in which  $A^3$  is phenyl or heteroaryl containing a total of 5 to 9 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-R^9$ ,  $-X^2OR^9$ ,  $-X^2SR^9$  and  $-X^2S(O)_2R^9$ , wherein  $R^9$  is  $-X^2R^{10}$ ,  $X^2$  is a bond or  $(C_{1-6})$ alkylene and  $R^{10}$  is phenyl or heteroaryl containing a total of 5 to 6 ring atoms, wherein each ring within  $A^3$  and  $R^{10}$  may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, halo, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)_2R^6$  and  $-X^2NR^4R^4$ , wherein  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl; and the *N*-oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

54. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Claim 1, 47 or 51 or a *N*-oxide derivative, prodrug derivative, individual isomer or mixture of isomers or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable excipient.

55. The pharmaceutical composition of Claim 54, further comprising at least one known cancer chemotherapeutic agent.

56. The pharmaceutical composition of Claim 55, wherein said cancer therapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C; carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguanzone, epirubicin, aclarubicin, bleomycin, imitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin®, Rituxan® and alanosine.

57. A method of treating a disorder responsive to the induction of apoptosis in an animal suffering said disorder, comprising administering to a mammal in need of such treatment an effective amount of a compound of Formula I:



in which:

the dashed lines indicate optional unsaturation without violating valency rules;

$R^1$  is hydrogen,  $(C_{1-6})$ alkyl or  $-C(O)R^6$ , wherein  $R^6$  is as defined below, or  $R^1$  is absent when a double bond exists between the nitrogen atom to which  $R^1$  is attached and an adjacent ring atom or  $R^1$  is as defined below;

$X^1$  is  $-NR^2$ -,  $-S$ -,  $-S(O)$ -,  $-S(O)_2$ - or  $-O$ -, wherein  $R^2$  is hydrogen or  $(C_{1-6})$ alkyl or is absent when a double bond exists between the nitrogen atom to which  $R^2$  is attached and an adjacent ring atom;

$A^1$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, or  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a fused polycyclic ring system selected from heteroaryl and unsaturated, partially unsaturated or saturated heterocycloalkyl in any case containing a total of 10 to 15 ring atoms, wherein  $A^1$  may be substituted with a group selected from  $-R^3$ ,  $-X^2OR^3$ ,  $-X^2C(O)R^3$ ,  $-X^2OC(O)R^3$ ,  $-X^2C(O)OR^3$ ,  $-X^2SR^3$ ,  $-X^2S(O)R^3$ ,  $-X^2S(O)_2R^3$ ,  $-X^2NR^3R^4$ ,  $-X^2NR^4C(O)R^3$ ,  $-X^2NR^4C(O)OR^3$ ,  $-X^2C(O)NR^3R^4$ ,  $-X^2NR^4C(O)NR^3R^4$ ,  $-X^2NR^4C(NR^4)NR^3R^4$ ,  $-X^2NR^4S(O)_2R^3$  and  $-X^2S(O)_2NR^3R^4$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^3$  is  $-X^2R^5$  wherein  $X^2$  is as defined above and  $R^5$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^1$  and  $R^5$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within  $A^1$  and  $R^5$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the provisos that only one of  $A^1$  and  $R^5$  is a fused polycyclic ring system;

$A^2$  is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein  $A^2$  may be substituted with a group selected




from  $-R^8$ ,  $-X^2OR^8$ ,  $-X^2C(O)R^8$ ,  $-X^2OC(O)R^8$ ,  $-X^2C(O)OR^8$ ,  $-X^2SR^8$ ,  $-X^2S(O)R^8$ ,  $-X^2S(O)_2R^8$ ,  $-X^2NR^4R^8$ ,  $-X^2NR^4C(O)R^8$ ,  $-X^2NR^4C(O)OR^8$ ,  $-X^2C(O)NR^4R^8$ ,  $-X^2NR^4C(O)NR^4R^8$ ,  $-X^2NR^4C(NR^4)NR^4R^8$ ,  $-X^2NR^4S(O)_2R^8$  and  $-X^2S(O)_2NR^4R^8$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$  is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^2$  and  $R^8$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2C(O)NR^4X^2C(O)OR^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said heterocycloalkylene, carbocycloalkyl and heterocycloalkyl rings within  $A^2$  and  $R^8$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^2$  and  $R^8$  is a fused polycyclic ring system; and

$A^3$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-R^9$ ,  $-X^2OR^9$ ,  $-X^2C(O)R^9$ ,  $-X^2OC(O)R^9$ ,  $-X^2C(O)OR^9$ ,  $-X^2SR^9$ ,  $-X^2S(O)R^9$ ,  $-X^2S(O)_2R^9$ ,  $-X^2NR^4R^9$ ,  $-X^2NR^4C(O)R^9$ ,  $-X^2NR^4C(O)OR^9$ ,  $-X^2C(O)NR^4R^9$ ,  $-X^2NR^4C(O)NR^4R^9$ ,  $-X^2NR^4C(NR^4)NR^4R^9$ ,  $-X^2NR^4S(O)_2R^9$  and  $-X^2S(O)_2NR^4R^9$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^9$  is  $-X^2R^{10}$  wherein  $X^2$  is as defined above and  $R^{10}$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated,

-125-

partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^3$  and  $R^{10}$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within  $A^3$  and  $R^{10}$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the proviso that only one of  $A^3$  and  $R^{10}$  is a fused polycyclic ring system; or an *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers, or a pharmaceutically acceptable salt thereof; with the proviso that when said compound is of Formula II(a):




II(a)

then  $A^3$  is other than:

- (a) benzo[1,3]dioxolyl;
- (b) phenyl which is mono-substituted by bromo, hydroxy, methyl or isopropyl; and
- (c) phenyl which is substituted by at least one of Cl and methoxy and not substituted by methylsulfanyl, amino, methylamino and dimethylamino.

58. The method of claim 57, with the further proviso that when said compound is selected from the group consisting of Formula II(a) and II(b):


-126-



then A<sup>3</sup> is other than:

- (a) benzo[1,3]dioxolyl;
- (b) phenyl which is mono-substituted by bromo, nitro, hydroxy, methyl, or isopropyl; and
- (c) phenyl which is substituted by at least one of Cl and methoxy and not substituted by methylsulfanyl, amino, methylamino and dimethylamino.

59. The method of claim 57, with the further proviso that when said compound is selected from the group consisting of Formula II(a) and II(b):




then A<sup>3</sup> is other than:


- (a) benzo[1,3]dioxolyl;
- (b) 2,3-dihydro-benzo[1,4]dioxinyl; and
- (c) phenyl which is substituted by at least one of bromo, chloro, hydroxy, nitro, methoxy and (C<sub>1-3</sub>)alkyl.

-127-


60. The method of Claim 57, wherein  $A^1$  of said compound is a group selected from Formulae (a), (b), (c), (d) and (e):




(a)




(b)



(c)




(d)



(e)

in which  $R^7$  is hydrogen or methyl,  $R^{11}$  is hydrogen or  $(C_{1-6})$ alkyl and the free valance is attached to  $A^2$ , or


$A^2$  and  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a group of Formula (g):




(g)

wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ ; and


$A^2$  of said compound is as defined above or is a group selected from Formulae (h), (i), (j), (k), (l) and (m):




(h)




(i)




(j)



(k)



(l)



(m)

in which  $n$  is 1 or 2 and  $R^1$  is acetyl or trifluoroacetyl; or an  $N$ -oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers, or a pharmaceutically acceptable salt thereof.

61. The method of Claim 60, wherein  $A^3$  of said compound is phenyl or heteroaryl containing a total of 5 to 9 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-R^9$ ,  $-X^2OR^9$ ,  $-X^2SR^9$  and  $-X^2S(O)_2R^9$ , wherein  $R^9$  is  $-X^2R^{10}$ ,  $X^2$  is a bond or  $(C_{1-6})$ alkylene and  $R^{10}$  is phenyl or heteroaryl containing a total of 5 to 6 ring atoms, wherein each ring within  $A^3$  and  $R^{10}$  may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, halo, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)_2R^6$  and  $-X^2NR^4R^4$ , wherein  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl; and the  $N$ -oxide derivatives, prodrug derivatives, protected derivatives, individual stereoisomers and mixtures of stereoisomers; and the pharmaceutically acceptable salts thereof.

62. The method of Claim 61, wherein said compound is selected from the group consisting of:

2-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

5 4-hydroxy-3-[7-(4-methanesulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

2-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

15 3-hydroxy-2-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-cyclohex-2-enone;

3-hydroxy-2-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

20 4-hydroxy-6-methyl-3-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-5,6-dihydro-pyran-2-one; and

4-hydroxy-6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-5,6-dihydro-pyran-2-one; or

25 a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

63. The method of claim 57, wherein said compound is selected from the list consisting of:

4-hydroxy-6-methyl-3-[7-(4-methylsulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[4-acetyl-7-(2,4-dimethoxy-phenyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,4-dimethoxy-phenyl)-4-(2,2,2-trifluoro-ethanoyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5           1-[7-(2,4-dimethoxy-phenyl)-5-(3-fluoro-4-methoxyphenyl)-[1,4]thiazepan-4-yl]-ethanone;

4-hydroxy-6-methyl-3-[7-(3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

10           3-[7-(5-ethyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(1-benzyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(2-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

15           4-hydroxy-6-methyl-3-[7-(3-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(4-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

20           4-hydroxy-6-methyl-3-[7-[3-(3-trifluoromethyl-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-[3-(3,4-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

25           4-hydroxy-6-methyl-3-{7-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

3-{7-[5-(2-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

30           3-{7-[5-(3-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[5-(4-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-{7-[5-(2-chloro-5-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

5           3-[7-(4-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

10           3-[7-(1-benzenesulfonyl-1*H*-pyrrol-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(3-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

15           4-hydroxy-6-methyl-3-[7-(1-methyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(3-chloro-2-methyl-5-trifluoromethyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

20           3-{7-[1-(2,4-difluoro-benzenesulfonyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

25           3-{7-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-chloro-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

30           4-hydroxy-6-methyl-3-[7-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;



3-[7-(4,5-dibromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2-chloro-5-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methylsulfanyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

10 3-[7-(5-chloro-1-methyl-3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[4-(2,4-dimethoxy-phenyl)-4,5-dihydro-3*H*-benzo[*b*][1,4]diazepin-2-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

15 3-[7-2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-1*H*-quinolin-2-one;

4-hydroxy-6-methyl-3-[7-(4-trifluoromethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

20 3-[7-(bis-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

25 3-hydroxy-2-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

3-[7-(2,4-dimethoxy-phenyl)-1-oxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>4</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methoxy-pyran-2-one;

10-(2,4-dimethoxy-phenyl)-3-methyl-7,8-dihydro-10*H*-2,5-dioxa-9-thia-6*a*-aza-cyclohepta[*a*]naphthalene-1,6-dione;

30 3-[7-(2,4-dimethoxy-phenyl)-1,1-dioxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>6</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5           2-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone; and

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one; or

10           a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

64.     A method of treating a disorder responsive to the induction of apoptosis in an animal suffering said disorder, comprising administering to a mammal in need of such treatment an effective amount of a compound selected from the group consisting of:

15           4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(4-ethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

20           3-[7-(3-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[7-(2-bromo-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

25           3-[7-(2,3-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

6-methyl-3-(2-*p*-tolyl-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl)-pyran-2-one;

4-hydroxy-6-methyl-3-[2-(4-methylsulfanyl-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-pyran-2-one; and

5 3-[7-(2,4-dimethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-3-[7-(4-chloro-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; and

10 4-hydroxy-3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; or

a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

15 65. The method of claim 64, wherein said compound is selected from the group consisting of:

4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;


3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one; and

20 6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one; or

a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

25 66. A method for treating or preventing cancer, comprising administering to an animal in need of such treatment an effective amount of a compound of Formula I:

-135-



in which:

the dashed lines indicate optional unsaturation without violating valency rules;

$R^1$  is hydrogen,  $(C_{1-6})$ alkyl or  $-C(O)R^6$ , wherein  $R^6$  is as defined below, or  $R^1$  is absent when a double bond exists between the nitrogen atom to which  $R^1$  is attached and an adjacent ring atom or  $R^1$  is as defined below;

$X^1$  is  $-NR^2-$ ,  $-S-$ ,  $-S(O)-$ ,  $-S(O)_2-$  or  $-O-$ , wherein  $R^2$  is hydrogen or  $(C_{1-6})$ alkyl or is absent when a double bond exists between the nitrogen atom to which  $R^2$  is attached and an adjacent ring atom;

$A^1$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, or  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a fused polycyclic ring system selected from heteroaryl and unsaturated, partially unsaturated or saturated heterocycloalkyl in any case containing a total of 10 to 15 ring atoms, wherein  $A^1$  may be substituted with a group selected from  $-R^3$ ,  $-X^2OR^3$ ,  $-X^2C(O)R^3$ ,  $-X^2OC(O)R^3$ ,  $-X^2C(O)OR^3$ ,  $-X^2SR^3$ ,  $-X^2S(O)R^3$ ,  $-X^2S(O)_2R^3$ ,  $-X^2NR^3R^4$ ,  $-X^2NR^4C(O)R^3$ ,  $-X^2NR^4C(O)OR^3$ ,  $-X^2C(O)NR^3R^4$ ,  $-X^2NR^4C(O)NR^3R^4$ ,  $-X^2NR^4C(NR^4)NR^3R^4$ ,  $-X^2NR^4S(O)_2R^3$  and  $-X^2S(O)_2NR^3R^4$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^3$  is  $-X^2R^5$  wherein  $X^2$  is as defined above and  $R^5$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^1$  and  $R^5$  contains from 3 to 8 ring atoms and may be


substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within A<sup>1</sup> and R<sup>5</sup> may be substituted further with 1 to 2 groups independently selected from (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo, with the provisos that only one of A<sup>1</sup> and R<sup>5</sup> is a fused polycyclic ring system;

A<sup>2</sup> is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein A<sup>2</sup> may be substituted with a group selected from -R<sup>8</sup>, -X<sup>2</sup>OR<sup>8</sup>, -X<sup>2</sup>C(O)R<sup>8</sup>, -X<sup>2</sup>OC(O)R<sup>8</sup>, -X<sup>2</sup>C(O)OR<sup>8</sup>, -X<sup>2</sup>SR<sup>8</sup>, -X<sup>2</sup>S(O)R<sup>8</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>8</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>8</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>8</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>8</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>8</sup> is -X<sup>2</sup>R<sup>9</sup> wherein X<sup>2</sup> is as defined above and R<sup>9</sup> is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, wherein each ring within A<sup>2</sup> and R<sup>8</sup> contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said heterocycloalkylene, carbocycloalkyl and heterocycloalkyl rings within A<sup>2</sup> and R<sup>8</sup> may be substituted further with 1 to 2 groups independently selected from

(C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo, with the proviso that only one of A<sup>2</sup> and R<sup>8</sup> is a fused polycyclic ring system; and

A<sup>3</sup> is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14  
5 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, wherein A<sup>3</sup> may be substituted with a group selected from -R<sup>9</sup>, -X<sup>2</sup>OR<sup>9</sup>, -X<sup>2</sup>C(O)R<sup>9</sup>, -X<sup>2</sup>OC(O)R<sup>9</sup>, -X<sup>2</sup>C(O)OR<sup>9</sup>, -X<sup>2</sup>SR<sup>9</sup>, -X<sup>2</sup>S(O)R<sup>9</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>9</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>9</sup>,  
10 -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>9</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>9</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>9</sup> is -X<sup>2</sup>R<sup>10</sup> wherein X<sup>2</sup> is as defined above and R<sup>10</sup> is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and R<sup>4</sup> at each occurrence independently  
15 is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, wherein each ring within A<sup>3</sup> and R<sup>10</sup> contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>,  
20 -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within A<sup>3</sup> and R<sup>10</sup> may be substituted further with 1 to 2  
25 groups independently selected from (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo, with the proviso that only one of A<sup>3</sup> and R<sup>10</sup> is a fused polycyclic ring system; or an N-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers, or a pharmaceutically acceptable salt thereof; with the proviso that when said compound is of Formula II(a):

-138-




II(a)

then  $A^3$  is other than:

- (a) benzo[1,3]dioxolyl;
- (b) phenyl which is mono-substituted by bromo, hydroxy, methyl or isopropyl; and
- (c) phenyl which is substituted by at least one of Cl and methoxy and not substituted by methylsulfanyl, amino, methylamino and dimethylamino; or a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

67. The method of claim 66, with the further proviso that when said compound is selected the group consisting of Formula II(a) and II(b):



II(a)


II(b)

then  $A^3$  is other than:

- (a) benzo[1,3]dioxolyl;
- (b) phenyl which is mono-substituted by bromo, nitro, hydroxy, methyl, or isopropyl; and
- (c) phenyl which is substituted by at least one of Cl and methoxy and not substituted by methylsulfanyl, amino, methylamino and dimethylamino; or

a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

68. The method of claim 66, with the further proviso that when said compound is selected the group consisting of Formula II(a) and II(b):



then A<sup>3</sup> is other than:


- (a) benzo[1,3]dioxolyl;
- (b) 2,3-dihydro-benzo[1,4]dioxinyl; and
- (c) phenyl which is substituted by at least one of bromo, chloro, hydroxy, nitro, methoxy and (C<sub>1-3</sub>)alkyl; or

a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof..


69. The method of Claim 66, wherein A<sup>1</sup> of said compound is a group selected from Formulae (a), (b), (c), (d) and (e):




-140-




(a)




(b)



(c)



(d)




(e)

in which  $R^7$  is hydrogen or methyl,  $R^{11}$  is hydrogen or  $(C_{1-6})$ alkyl and the free valance is attached to  $A^2$ , or

$A^2$  and  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a group of Formula (g):

5




(g)

wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ ; and

$A^2$  of said compound is as defined above or is a group selected from Formulae (h), (i), (j), (k), (l) and (m):

-141-



in which  $n$  is 1 or 2 and  $R^1$  is acetyl or trifluoroacetyl; or a  $N$ -oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

- 5                      70.    The method of Claim 69, wherein  $A^3$  of said compound is phenyl or heteroaryl containing a total of 5 to 9 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-R^9$ ,  $-X^2OR^9$ ,  $-X^2SR^9$  and  $-X^2S(O)_2R^9$ , wherein  $R^9$  is  $-X^2R^{10}$ ,  $X^2$  is a bond or  $(C_{1-6})$ alkylene and  $R^{10}$  is phenyl or heteroaryl containing a total of 5 to 6 ring atoms, wherein each ring within  $A^3$  and  $R^{10}$  may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, halo, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)_2R^6$  and  $-X^2NR^4R^4$ , wherein  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl; or a  $N$ -oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.
- 10
- 15

71.    The method of Claim 70, wherein said compound is selected from the group consisting of:

4-hydroxy-3-[7-(2-methoxy-4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

2-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

5 4-hydroxy-3-[7-(4-methanesulfonyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

2-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone;

15 3-hydroxy-2-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-cyclohex-2-enone;

3-hydroxy-2-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

20 4-hydroxy-6-methyl-3-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-5,6-dihydro-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-5,6-dihydro-pyran-2-one; and

3-[7-(2,4-dimethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one; or

25 a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

72. The method of claim 66, wherein said compound is selected from the list consisting of:

3-[4-acetyl-7-(2,4-dimethoxy-phenyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,4-dimethoxy-phenyl)-4-(2,2,2-trifluoro-ethanoyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 1-[7-(2,4-dimethoxy-phenyl)-5-(3-fluoro-4-methoxyphenyl)-[1,4]thiazepan-4-yl]-ethanone;

4-hydroxy-6-methyl-3-[7-(3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

10 3-[7-(5-ethyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(1-benzyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3[7-(2-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

15 4-hydroxy-6-methyl-3[7-(3-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3[7-(4-trifluoromethylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

20 4-hydroxy-6-methyl-3-[7-[3-(3-trifluoromethyl-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-[3-(3,4-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

25 4-hydroxy-6-methyl-3-{7-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

3-{7-[5-(2-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

30 3-{7-[5-(3-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[5-(4-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-{7-[5-(2-chloro-5-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

5 3-[7-(4-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 3-[7-(1-benzenesulfonyl-1*H*-pyrrol-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(3-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

15 4-hydroxy-6-methyl-3-[7-(1-methyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(3-chloro-2-methyl-5-trifluoromethyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

20 3-{7-[1-(2,4-difluoro-benzenesulfonyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

25 3-{7-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-chloro-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

30 4-hydroxy-6-methyl-3-[7-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(4,5-dibromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2-chloro-5-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methylsulfanyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

10 3-[7-(5-chloro-1-methyl-3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[4-(2,4-dimethoxy-phenyl)-4,5-dihydro-3*H*-benzo[*b*][1,4]diazepin-2-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

15 3-[7-2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-1*H*-quinolin-2-one;

4-hydroxy-6-methyl-3-[7-(4-trifluoromethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

20 3-[7-(bis-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-hydroxy-2-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

25 3-[7-(2,4-dimethoxy-phenyl)-1-oxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>4</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methoxy-pyran-2-one;

10-(2,4-dimethoxy-phenyl)-3-methyl-7,8-dihydro-10*H*-2,5-dioxa-9-thia-6a-aza-cyclohepta[*a*]naphthalene-1,6-dione;

30 3-[7-(2,4-dimethoxy-phenyl)-1,1-dioxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>6</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl)-2,3-dihydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one;

5 2-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone; and

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one; or

10 a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

73. A method for treating or preventing cancer, comprising administering to an animal in need of such treatment an effective amount of a compound selected from the group consisting of:

15 4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(4-ethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

20 3-[7-(3-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[7-(2-bromo-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,3-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

25 3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

30 6-methyl-3-(2-*p*-tolyl-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl)-pyran-2-one;

4-hydroxy-6-methyl-3-[2-(4-methylsulfanyl-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-pyran-2-one;

4-hydroxy-3-[7-(4-chloro-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; and

5 4-hydroxy-3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; or

a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

10 74. The method of claim 73, wherein said compound is selected from the group consisting of:

4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

15 3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one; and

6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one; or


20 a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

25 75. The method of Claims 66 and 73, wherein said cancer is selected from the group consisting of Hodgkin's disease, non-Hodgkin's lymphoma, acute and chronic lymphocytic leukemias, multiple myeloma, neuroblastoma, breast carcinoma, ovarian carcinoma, lung carcinoma, Wilms' tumor, cervical carcinoma, testicular carcinoma, soft-tissue sarcoma, chronic lymphocytic leukemia, primary macroglobulinemia, bladder carcinoma, chronic granulocytic leukemia, primary brain carcinoma, malignant melanoma, small-cell lung carcinoma, stomach carcinoma, colon carcinoma, malignant pancreatic insulinoma, malignant carcinoid carcinoma, choriocarcinoma, mycosis fungoides,



head and neck carcinoma, osteogenic sarcoma, pancreatic carcinoma, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinoma, thyroid carcinoma, esophageal carcinoma, malignant hypercalcemia, cervical hyperplasia, renal cell carcinoma, endometrial carcinoma, polycythemia vera, essential thrombocytosis, adrenal cortex carcinoma, skin cancer and prostatic carcinoma.

76. A method for the treatment of drug resistant cancer, comprising administering to an animal in need of such treatment an effective amount of a compound of compound of Formula I:



in which:

the dashed lines indicate optional unsaturation without violating valency rules;

$R^1$  is hydrogen,  $(C_{1-6})$ alkyl or  $-C(O)R^6$ , wherein  $R^6$  is as defined below, or  $R^1$  is absent when a double bond exists between the nitrogen atom to which  $R^1$  is attached and an adjacent ring atom or  $R^1$  is as defined below;

$X^1$  is  $-NR^2$ -,  $-S$ -,  $-S(O)$ -,  $-S(O)_2$ - or  $-O$ -, wherein  $R^2$  is hydrogen or  $(C_{1-6})$ alkyl or is absent when a double bond exists between the nitrogen atom to which  $R^2$  is attached and an adjacent ring atom;

$A^1$  is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, or  $A^1$  together with  $R^1$  and the atoms to which  $A^1$  and  $R^1$  are attached forms a fused polycyclic ring system selected from heteroaryl and unsaturated, partially unsaturated or saturated heterocycloalkyl in any case containing a total of 10 to 15 ring atoms, wherein  $A^1$

may be substituted with a group selected from  $-R^3$ ,  $-X^2OR^3$ ,  $-X^2C(O)R^3$ ,  $-X^2OC(O)R^3$ ,  $-X^2C(O)OR^3$ ,  $-X^2SR^3$ ,  $-X^2S(O)R^3$ ,  $-X^2S(O)_2R^3$ ,  $-X^2NR^3R^4$ ,  $-X^2NR^4C(O)R^3$ ,  $-X^2NR^4C(O)OR^3$ ,  $-X^2C(O)NR^3R^4$ ,  $-X^2NR^4C(O)NR^3R^4$ ,  $-X^2NR^4C(NR^4)NR^3R^4$ ,  $-X^2NR^4S(O)_2R^3$  and  $-X^2S(O)_2NR^3R^4$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^3$  is  $-X^2R^5$  wherein  $X^2$  is as defined above and  $R^5$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, wherein each ring within  $A^1$  and  $R^5$  contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from  $(C_{1-6})$ alkyl, cyano, halo, nitro, halo-substituted  $(C_{1-6})$ alkyl,  $-X^2OR^4$ ,  $-X^2C(O)R^6$ ,  $-X^2OC(O)R^6$ ,  $-X^2C(O)OR^4$ ,  $-X^2SR^4$ ,  $-X^2S(O)R^6$ ,  $-X^2S(O)_2R^6$ ,  $-X^2NR^4R^4$ ,  $-X^2NR^4C(O)R^6$ ,  $-X^2NR^4C(O)OR^4$ ,  $-X^2C(O)NR^4R^4$ ,  $-X^2NR^4C(O)NR^4R^4$ ,  $-X^2NR^4C(NR^4)NR^4R^4$ ,  $-X^2NR^4S(O)_2R^6$  and  $-X^2S(O)_2NR^4R^4$ , wherein  $X^2$  and  $R^4$  are as defined above and  $R^6$  is  $(C_{1-6})$ alkyl or halo-substituted  $(C_{1-6})$ alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within  $A^1$  and  $R^5$  may be substituted further with 1 to 2 groups independently selected from  $(C_{1-6})$ alkylidene, oxo, imino and thioxo, with the provisos that only one of  $A^1$  and  $R^5$  is a fused polycyclic ring system;


$A^2$  is a monocyclic or fused bicyclic ring selected from heteroarylene or unsaturated, partially unsaturated or saturated heterocycloalkylene containing a total of 5 to 11 ring atoms, wherein  $A^2$  may be substituted with a group selected from  $-R^8$ ,  $-X^2OR^8$ ,  $-X^2C(O)R^8$ ,  $-X^2OC(O)R^8$ ,  $-X^2C(O)OR^8$ ,  $-X^2SR^8$ ,  $-X^2S(O)R^8$ ,  $-X^2S(O)_2R^8$ ,  $-X^2NR^4R^8$ ,  $-X^2NR^4C(O)R^8$ ,  $-X^2NR^4C(O)OR^8$ ,  $-X^2C(O)NR^4R^8$ ,  $-X^2NR^4C(O)NR^4R^8$ ,  $-X^2NR^4C(NR^4)NR^4R^8$ ,  $-X^2NR^4S(O)_2R^8$  and  $-X^2S(O)_2NR^4R^8$ , wherein  $X^2$  is a bond or  $(C_{1-6})$ alkylene,  $R^8$  is  $-X^2R^9$  wherein  $X^2$  is as defined above and  $R^9$  is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and  $R^4$  at each occurrence independently is hydrogen,  $(C_{1-6})$ alkyl or

halo-substituted (C<sub>1-6</sub>)alkyl, wherein each ring within A<sup>2</sup> and R<sup>8</sup> contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said heterocycloalkylene, carbocycloalkyl and heterocycloalkyl rings within A<sup>2</sup> and R<sup>8</sup> may be substituted further with 1 to 2 groups independently selected from (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo, with the proviso that only one of A<sup>2</sup> and R<sup>8</sup> is a fused polycyclic ring system; and

A<sup>3</sup> is a monocyclic or fused polycyclic ring system selected from aryl containing a total of 6 to 14 ring atoms, heteroaryl containing a total of 5 to 14 ring atoms and unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 14 ring atoms, wherein A<sup>3</sup> may be substituted with a group selected from -R<sup>9</sup>, -X<sup>2</sup>OR<sup>9</sup>, -X<sup>2</sup>C(O)R<sup>9</sup>, -X<sup>2</sup>OC(O)R<sup>9</sup>, -X<sup>2</sup>C(O)OR<sup>9</sup>, -X<sup>2</sup>SR<sup>9</sup>, -X<sup>2</sup>S(O)R<sup>9</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>9</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>9</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>9</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>9</sup>, wherein X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene, R<sup>9</sup> is -X<sup>2</sup>R<sup>10</sup> wherein X<sup>2</sup> is as defined above and R<sup>10</sup> is aryl containing a total of 6 to 10 ring atoms, heteroaryl containing a total of 5 to 10 ring atoms or unsaturated, partially unsaturated or saturated carbocycloalkyl or heterocycloalkyl each containing a total of 3 to 10 ring atoms, and R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl, wherein each ring within A<sup>3</sup> and R<sup>10</sup> contains from 3 to 8 ring atoms and may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, cyano, halo, nitro, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>OC(O)R<sup>6</sup>, -X<sup>2</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)R<sup>6</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)R<sup>6</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)OR<sup>4</sup>, -X<sup>2</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(O)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>C(NR<sup>4</sup>)NR<sup>4</sup>R<sup>4</sup>, -X<sup>2</sup>NR<sup>4</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>S(O)<sub>2</sub>NR<sup>4</sup>R<sup>4</sup>, wherein X<sup>2</sup> and R<sup>4</sup> are as defined above and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or

-151-

halo-substituted (C<sub>1-6</sub>)alkyl, and wherein any said carbocycloalkyl and heterocycloalkyl rings within A<sup>3</sup> and R<sup>10</sup> may be substituted further with 1 to 2 groups independently selected from (C<sub>1-6</sub>)alkylidene, oxo, imino and thioxo, with the proviso that only one of A<sup>3</sup> and R<sup>10</sup> is a fused polycyclic ring system; or a  
 5 *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.; with the proviso that when said compound is of Formula II(a):




II(a)

then A<sup>3</sup> is other than:

- 10 (a) benzo[1,3]dioxolyl;  
 (b) phenyl which is mono-substituted by bromo, hydroxy, methyl or isopropyl; and  
 (c) phenyl which is substituted by at least one of Cl and methoxy and not substituted by methylsulfanyl, amino, methylamino and dimethylamino; or  
 15 a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

77. The method of claim 76, with the further proviso that when said compound is selected the group consisting of Formula II(a) and II(b):


-152-



then A<sup>3</sup> is other than:

- (a) benzo[1,3]dioxolyl;
- (b) phenyl which is mono-substituted by bromo, nitro, hydroxy,  
methyl or isopropyl; and
- (c) phenyl which is substituted by at least one of Cl and methoxy and  
not substituted by methylsulfanyl, amino, methylamino and dimethylamino; or  
a *N*-oxide derivative, prodrug derivative, protected derivative, individual  
stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt  
thereof.

78. The method of claim 76, with the further proviso that when said compound is selected the group consisting of Formula II(a) and II(b):



then A<sup>3</sup> is other than:


- (a) benzo[1,3]dioxolyl;
- (b) 2,3-dihydro-benzo[1,4]dioxinyl; and

(c) phenyl which is substituted by at least one of bromo, chloro, hydroxy, nitro, methoxy and (C<sub>1-3</sub>)alkyl; or


a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof. 1.

5


79. The method of Claim 76, wherein A<sup>1</sup> of said compound is a group selected from Formulae (a), (b), (c), (d) and (e):




(a)




(b)



(c)



(d)




(e)

in which R<sup>7</sup> is hydrogen or methyl, R<sup>11</sup> is hydrogen or (C<sub>1-6</sub>)alkyl and the free valance is attached to A<sup>2</sup>, or

10

A<sup>2</sup> and A<sup>1</sup> together with R<sup>1</sup> and the atoms to which A<sup>1</sup> and R<sup>1</sup> are attached forms a group of Formula (g):

-154-




(g)


wherein  $X^1$  is -S- and the free valance is attached to  $A^3$ ; and

$A^2$  of said compound is as defined above or is a group selected from Formulae (h), (i), (j), (k), (l) and (m):


5




(h)




(i)




(j)



(k)



(l)



(m)

in which  $n$  is 1 or 2 and  $R^1$  is acetyl or trifluoroacetyl; or a  $N$ -oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

10

80. The method of Claim 79, wherein  $A^3$  of said compound is phenyl or heteroaryl containing a total of 5 to 9 ring atoms, wherein  $A^3$  may be substituted with a group selected from  $-R^9$ ,  $-X^2OR^9$ ,  $-X^2SR^9$  and  $-X^2S(O)_2R^9$ ,

wherein R<sup>9</sup> is -X<sup>2</sup>R<sup>10</sup>, X<sup>2</sup> is a bond or (C<sub>1-6</sub>)alkylene and R<sup>10</sup> is phenyl or heteroaryl containing a total of 5 to 6 ring atoms, wherein each ring within A<sup>3</sup> and R<sup>10</sup> may be substituted with 1 to 3 groups independently selected from (C<sub>1-6</sub>)alkyl, halo, halo-substituted (C<sub>1-6</sub>)alkyl, -X<sup>2</sup>OR<sup>4</sup>, -X<sup>2</sup>SR<sup>4</sup>, -X<sup>2</sup>S(O)<sub>2</sub>R<sup>6</sup> and -X<sup>2</sup>NR<sup>4</sup>R<sup>4</sup>, wherein  
5 R<sup>4</sup> at each occurrence independently is hydrogen, (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl and R<sup>6</sup> is (C<sub>1-6</sub>)alkyl or halo-substituted (C<sub>1-6</sub>)alkyl; or a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

81. The method of Claim 80, wherein said compound is selected from  
10 the group consisting of:

4-hydroxy-3-[7-(2-methoxy-4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-  
[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

2-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-  
3-hydroxy-cyclohex-2-enone;

15 4-hydroxy-3-[7-(4-methanesulfonyl-phenyl)-2,3,6,7-tetrahydro-  
[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-  
4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-  
20 4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-1,4]thiazepin-5-yl]-  
4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

2-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-  
3-hydroxy-cyclohex-2-enone;

3-hydroxy-2-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-  
[1,4]thiazepin-5-yl]-cyclohex-2-enone;

3-hydroxy-2-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-  
benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

4-hydroxy-6-methyl-3-[2-(2,3,4-trimethoxy-phenyl)-2,3-dihydro-  
30 benzo[*b*][1,4]thiazepin-4-yl]-5,6-dihydro-pyran-2-one;



4-hydroxy-6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-  
2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-5,6-dihydro-pyran-2-one; and

3-[7-(2,4-dimethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-  
6-methyl-pyran-2-one; or

5 a *N*-oxide derivative, prodrug derivative, protected derivative, individual  
stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt  
thereof.

82. The method of claim 76, wherein said compound is selected from  
the list consisting of:

10 3-[4-acetyl-7-(2,4-dimethoxy-phenyl)-[1,4]thiazepan-5-yl]-4-hydroxy-6-  
methyl-pyran-2-one;

3-[7-(2,4-dimethoxy-phenyl)-4-(2,2,2-trifluoro-ethanoyl)-[1,4]thiazepan-  
5-yl]-4-hydroxy-6-methyl-pyran-2-one;

15 1-[7-(2,4-dimethoxy-phenyl)-5-(3-fluoro-4-methoxyphenyl)-  
[1,4]thiazepan-4-yl]-ethanone;

4-hydroxy-6-methyl-3-[7-(3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-  
[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(5-ethyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-  
hydroxy-6-methyl-pyran-2-one;

20 3-[7-(1-benzyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-  
hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3[7-(2-trifluoromethylsulfanyl-phenyl)-2,3,6,7-  
tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

25 4-hydroxy-6-methyl-3[7-(3-trifluoromethylsulfanyl-phenyl)-2,3,6,7-  
tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3[7-(4-trifluoromethylsulfanyl-phenyl)-2,3,6,7-  
tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-[3-(3-trifluoromethyl-phenoxy)-phenyl]-2,3,6,7-  
tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-[3-(3,4-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-[3-(3,5-dichloro-phenoxy)-phenyl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

5 4-hydroxy-6-methyl-3-{7-[5-(3-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl}-pyran-2-one;

3-{7-[5-(2-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 3-{7-[5-(3-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[5-(4-chloro-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-{7-[5-(2-chloro-5-trifluoromethyl-phenyl)-furan-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

15 3-[7-(4-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-bromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

20 3-[7-(1-benzenesulfonyl-1*H*-pyrrol-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(3-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

25 4-hydroxy-6-methyl-3-[7-(1-methyl-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(3-chloro-2-methyl-5-trifluoromethyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

30 3-{7-[1-(2,4-difluoro-benzenesulfonyl)-1*H*-pyrrol-2-yl]-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one;

3-{7-[1-(3,5-dichloro-phenyl)-1*H*-pyrrol-2-yl]}-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one;

5 3-{7-[1-(4-chloro-phenyl)-1*H*-pyrrol-2-yl]}-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl)-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(5-chloro-1*H*-indol-3-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

10 4-hydroxy-6-methyl-3-[7-(6-*p*-tolylsulfanyl-imidazo[2,1-*b*]thiazol-5-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(4,5-dibromo-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2-chloro-5-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

15 3-[7-(2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one;

4-hydroxy-6-methyl-3-[7-(5-methylsulfanyl-thien-2-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

20 3-[7-(5-chloro-1-methyl-3-phenyl-1*H*-pyrazol-4-yl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[4-(2,4-dimethoxy-phenyl)-4,5-dihydro-3*H*-benzo[*b*][1,4]diazepin-2-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

25 3-[7-2,4-dimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-1*H*-quinolin-2-one;

4-hydroxy-6-methyl-3-[7-(4-trifluoromethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

30 3-[7-(bis-trifluoromethyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(4-dimethylamino-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-hydroxy-2-[2-(2,4-diethoxy-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-cyclohex-2-enone;

5 3-[7-(2,4-dimethoxy-phenyl)-1-oxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>4</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methoxy-pyran-2-one;

10-(2,4-dimethoxy-phenyl)-3-methyl-7,8-dihydro-10*H*-2,5-dioxa-9-thia-6a-aza-cyclohepta[*a*]naphthalene-1,6-dione;

3-[7-(2,4-dimethoxy-phenyl)-1,1-dioxo-2,3,6,7-tetrahydro-1*H*-1λ<sup>6</sup>-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

3-(7-[2,2']bithienyl-5-yl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

15 2-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-3-hydroxy-cyclohex-2-enone; and

3-[7-(2,4-diethoxy-phenyl)-2,3-dihydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-5,6-dihydro-pyran-2-one; or

a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

83. A method for the treatment of drug resistant cancer, comprising administering to an animal in need of such treatment an effective amount of a compound selected from the group consisting of:

25 4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

3-[7-(4-ethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

30 3-[7-(3-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one;

3-[7-(2-bromo-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]4-hydroxy-6-methyl-pyran-2-one;

3-[7-(2,3-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]4-hydroxy-6-methyl-pyran-2-one;

5 3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one;

6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

10 6-methyl-3-(2-*p*-tolyl-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl)-pyran-2-one;

4-hydroxy-6-methyl-3-[2-(4-methylsulfanyl-phenyl)-2,3-dihydro-benzo[*b*][1,4]thiazepin-4-yl]-pyran-2-one;

4-hydroxy-3-[7-(4-chloro-2-methoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; and

15 4-hydroxy-3-[7-(2,4-diethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-6-methyl-pyran-2-one; or

a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

20 84. The method of claim 83, wherein said compound is selected from the group consisting of:

4-hydroxy-6-methyl-3-[7-(4-methylsulfanyl-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one;

25 3-[7-(3,4-dichloro-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-4-hydroxy-6-methyl-pyran-2-one; and

6-methyl-3-[7-(2,3,4-trimethoxy-phenyl)-2,3,6,7-tetrahydro-[1,4]thiazepin-5-yl]-pyran-2-one; or

30 a *N*-oxide derivative, prodrug derivative, protected derivative, individual stereoisomer or mixture of stereoisomers; or a pharmaceutically acceptable salt thereof.

85. The method of Claim 66, 73, 76 or 83, further comprising administering to said animal at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent.

5 86. The method of Claim 85, wherein said known cancer therapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 10 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin, imitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin, Rituxan and alanosine.

87. The method of Claim 66, 73, 75 or 83, further comprising treating said animal with radiation-therapy.

15 88. The method of Claim 66, 73, 76 or 83, wherein said compound is administered after surgical treatment for cancer.

89. The method of Claim 57, wherein said disorder is an autoimmune disease.

90. The method of Claim 57, wherein said disorder is rheumatoid arthritis.

20 91. The method of Claim 57, wherein said disorder is inflammation or inflammatory bowel disease.

92. The method of Claim 57, wherein said disorder is psoriasis.

93. The method of Claim 57, wherein said disorder is a skin disease.

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number  
**WO 01/79187 A3**

- (51) International Patent Classification<sup>7</sup>: C07D 281/06, 281/10, 417/04, 417/14, 513/04, 513/14, 405/04, A61K 31/55, A61P 35/00
- (74) Agents: ESMOND, Robert, W. et al.: Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).
- (21) International Application Number: PCT/US01/12581
- (22) International Filing Date: 18 April 2001 (18.04.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/197,599 18 April 2000 (18.04.2000) US
- (71) Applicant (for all designated States except US): CYTOVIA, INC. [US/US]: 6650 Nancy Ridge Drive, San Diego, CA 92121 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicants and
- (72) Inventors: CAI, Sui, Xiong [US/US]: 3623 Berryfield Court, San Diego, CA 92130 (US). DREWE, John, A. [US/US]: 7069 Murilla Lane, Carlsbad, CA 92009 (US). SHELTON, Emma, Jane [US/US]: 680 Lemon Street, Menlo Park, CA 94025 (US). LITVAK, Joane [US/US]: 555 Jean Street, Apartment 626, Oakland, CA 94610 (US). SPERANDIO, David [CH/US]: 150 Paseo Court, Mountain View, CA 94043 (US). SPENCER, Jeffrey, R. [US/US]: 8 Baycrest Way, South San Francisco, CA 94080 (US).
- Published:  
— with international search report
- (88) Date of publication of the international search report:  
21 February 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED 1,4-THIAZEPINE AND ANALOGS AND THEIR USE AS ACTIVATORS OF CASPASES

(57) Abstract: The present invention is directed to substituted 1,4-thiazepine and analogs thereof, represented by the general Formula (I): wherein the dashed lines, A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup>, X<sup>1</sup> and R<sup>1</sup> are defined herein. The present invention also relates to the discovery that compounds having Formula I are activators of caspases and inducers of apoptosis. Therefore, the activators of caspases and inducers of apoptosis of this invention can be used to induce cell death in a variety of clinical conditions in which uncontrolled growth and spread of abnormal cells occurs.

# INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/US 01/12581

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D281/06 C07D281/10 C07D417/04 C07D417/14 C07D513/04  
C07D513/14 C07D405/04 A61K31/55 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 07017 A (PRESIDENT AND FELLOWS OF HARVARD COLLEGE) 10 February 2000 (2000-02-10) the whole document, particularly fig. 24, compounds 2a-2f, and fig. 25, compounds 2g-2q	1-56
X	WO 98 56785 A (SUMITOMO PHARMACEUTICALS CO., LTD.) 17 December 1998 (1998-12-17) the whole document, particularly page 28, compound 16	1-56
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

10 October 2001

Date of mailing of the international search report

05/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Allard, M



# INTERNATIONAL SEARCH REPORT

Inter. l. onal Application No

PCT/US 01/12581

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FODILI M ET AL: "An efficient synthesis of new 2-pyronyl-1,5-benzodiazepine derivatives" SYNTHESIS, no. 5, May 1999 (1999-05), pages 811-814, XP002179780 the whole document	1-56
X	DE SARRO G ET AL: "5H-1,2,4-Oxadiazolo[5,4-d]1,5-benzothiazepines as anticonvulsant agents in DBA/2 mice" EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY, vol. 30, 1995, pages 925-929, XP004040221 the whole document	1-56
X	ATTIA A E-H ET AL: "Azachalcones. IV. Reactions of azachalcones with o-phenylenediamine" GAZZETTA CHIMICA ITALIANA, vol. 112, no. 9/10, 1982, pages 387-390, XP000985350 the whole document	1-56
X	SRIVASTAVA V K ET AL: "2-(2'-Hydroxyphenyl)-4-aryl-1,5-benzodiazepines as CNS active agents" ARZNEIMITTEL-FORSCHUNG, vol. 32(II), no. 12, 1982, pages 1512-1514, XP002179781 the whole document	1-56
X	CHEMICAL ABSTRACTS, vol. 99, no. 13, 26 September 1983 (1983-09-26) Columbus, Ohio, US; abstract no. 105221v, JADHAV K P ET AL: "Synthesis of 2,4-diaryl-2,3-dihydro-1,5-benzothiazepines and their 1,1-dioxides as antibacterial agents" page 584; XP002179782 abstract & INDIAN J. CHEM., SECT. B, vol. 22B, no. 2, 1983, pages 180-182, -/--	1-56

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/12581

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 91, no. 23, 3 December 1979 (1979-12-03). Columbus, Ohio, US; abstract no. 193222j, SAYED A A ET AL: "The behavior of some 3-substituted 4-hydroxy-1-alkyl (or phenyl) carbostyrils towards amines and hydrazines" page 638; XP002179783 abstract, compounds IV & EGYPT. J. CHEM., vol. 19, no. 5, 1976, pages 811-826,	1-53
X	CHEMICAL ABSTRACTS, vol. 124, no. 13, 25 March 1996 (1996-03-25) Columbus, Ohio, US; abstract no. 176050j, SUCHETA K ET AL: "Synthesis of some new 1,5-benzothiazepine derivatives" page 1291; XP002179784 abstract & INDIAN J. CHEM., SECT. B: ORG. CHEM. INCL. MED. CHEM., vol. 34B, no. 10, 1995, pages 893-894,	1-53
X	CHEMICAL ABSTRACTS, vol. 131, no. 7, 16 August 1999 (1999-08-16) Columbus, Ohio, US; abstract no. 87897t, IBRAHIM S S ET AL: "New quinolones and naphthyridinones bearing heterocyclic rings" page 667; XP002179785 abstract -& DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; Database accession no. 1999:315201 XP002179787 RN 229316-41-0, 229316-43-2 and 229316-45-4 & CHEM. PAP., vol. 53, no. 1, 1999, pages 53-64,	1-53

---  
-/--

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/12581

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>CHEMICAL ABSTRACTS, vol. 127, no. 3, 21 July 1997 (1997-07-21) Columbus, Ohio, US; abstract no. 34096s, IBRAHIM S S ET AL: "Synthesis of new 3-acryloyl-1,2-dihydro-4-hydroxy-1-methyl- 2-oxoquinoline derivatives and their behavior towards some nucleophiles" page 635; XP002179786 abstract -&amp; DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; Database accession no. 1997:293144 XP002179788 RN 190514-14-8, 190514-15-9, 190514-16-0, 190514-17-1, 190514-27-3, 190514-28-4 and 190514-37-5 abstract &amp; CHEM. PAP., vol. 51, no. 1, 1997, pages 33-42,</p>	1-53
A	<p>WO 99 56736 A (TRINITY COLLEGE DUBLIN) 11 November 1999 (1999-11-11) the whole document</p>	1,56

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-25, 27-57, 60-93 (all partly)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty of claims 1-6, 15, 22, 23, 26, 28, 35, 36, 41, 42 and 54-56. So many documents were retrieved that it is impossible to determine which parts of those claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT).

Furthermore, the scope of claims 1-25, 27-57 and 60-93, in as far as the expressions "prodrug derivatives" and "protected derivative" are concerned, is unclear (Article 6 PCT).

For these reasons it appears impossible to execute a meaningful search and/or to issue a complete search report over the whole breadth of the above mentioned claims.

The search and the report can only be considered complete for the subject-matter as recited in claims 7-14, 16-21, 24-25, 27, 29-34, 37-40, 43-53, 60-65, 69-74 and 79-84 in as far as no "prodrug derivatives" and/or "protected derivatives" are concerned.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/12581

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 0007017	A	10-02-2000	AU	5322399 A		21-02-2000
			WO	0007017 A2		10-02-2000
WO 9856785	A	17-12-1998	AU	7553198 A		30-12-1998
			WO	9856785 A1		17-12-1998
WO 9956736	A	11-11-1999	AU	3843899 A		23-11-1999
			WO	9956736 A2		11-11-1999

• • • • •

1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 2677, 2678, 2679, 26

100

• • • • •

•

•

•

•

•

•

• •

•

•  
•  
•

•  
•